arXiv:1011.1391v1 [math.NT] 5 Nov 2010

On Karatsuba's Problem Concerning the Divisor Function $\tau(n)$[1]

M.A. Korolev

Abstract. We study an asymptotic behavior of the sum $\sum_{n\leqslant x}\dfrac{\tau(n)}{\tau(n+a)}$. Here $\tau(n)$ denotes the number of divisors of n and $a\geqslant 1$ is a fixed integer.

1. Introduction

This paper deals with a problem stated by A.A. Karatsuba in November, 2004: to determine an asymptotic behavior of the following sum

$$S(x) \;=\; \sum_{n\leqslant x}\frac{\tau(n)}{\tau(n+1)}$$

(here $\tau(n)$ denotes the number of positive divisors of n). Since

$$\frac{1}{x}\sum_{n\leqslant x}\tau(n)\;\sim\;\ln x,\qquad \frac{1}{x}\sum_{n\leqslant x}\frac{1}{\tau(n)}\;\sim\;\frac{c}{\sqrt{\ln x}}$$

for some $c>0$, the below assumption seems reasonable:

$$\frac{1}{x}\,S(x)\;\sim\;\frac{K}{\sqrt{\ln x}}\cdot\ln x\;\sim\;K\sqrt{\ln x},\qquad K>0.$$

The aim of this paper is to prove the following

Theorem. *Let a be a fixed integer, $a\geqslant 1$, and let*

$$S_a(x)\;=\;\sum_{n\leqslant x}\frac{\tau(n)}{\tau(n+a)}.$$

Then

$$S_a(x)\;=\;K(a)x\sqrt{\ln x}\;+\;O(x\ln\ln x),$$

as $x\to+\infty$. The constant $K(a)$ has the form $K(a)=K\cdot\kappa(a)$, where

$$K=\frac{1}{\sqrt{\pi}}\prod_{p}\Bigl(\frac{1}{\sqrt{p(p-1)}}\;+\;\sqrt{1-\frac{1}{p}}(p-1)\ln\frac{p}{p-1}\Bigr)\;=\;0.757\,827\,651\ldots,$$

$$\kappa(a)=\beta(a)\prod_{p|a}\frac{1+\displaystyle\sum_{k=1}^{+\infty}\frac{e_a(p^k)}{k+1}\,p^{-k}}{1+\beta(p)\displaystyle\sum_{k=1}^{+\infty}\frac{p^{-k}}{k+1}},$$

$$e_a(n)=\frac{1}{\beta(a)}\sum_{d|(a,n)}\beta\Bigl(\frac{an}{d^2}\Bigr),\qquad \beta(a)=\prod_{p|a}\frac{(p-1)^2}{p^2-p+1}.$$

---

[1] This research was supported by the Programme of the President of the Russian Federation 'Young Candidates of the Russian Federation' (grant no. MK-4052.2009.1).

For the below, we need the following notations:

$\varphi(q)$ denotes Euler function;

χ denotes Dirichlet's character modulo q, $q \geqslant 3$;

the symbols $\sum\limits_{\substack{\chi \bmod q \\ \chi \neq \chi_0}}$ and $\sum\limits_{\chi \neq \chi_0}$ denote the sums over all non-principal characters modulo q;

the symbols $\sideset{}{^*}\sum\limits_{\chi \bmod q}$ and $\sideset{}{^*}\sum\limits_{\chi}$ denote the sums over all primitive characters modulo q;

$s = \sigma + it$, σ, t are real numbers;

$L(s, \chi)$ is Dirichlet's L-function corresponding to the character χ;

the symbol $N(\sigma; T, \chi)$ means the number of zeros of $L(s, \chi)$ in the rectangle $\sigma < \operatorname{Re} s \leqslant 1$, $|\operatorname{Im} s| \leqslant T$;

(a, b) denotes the great common divisor of a and b;

symbols $\theta, \theta_1, \theta_2, \ldots$ denote complex numbers such that $|\theta|,|\theta_1|,|\theta_2|,\ldots \leqslant 1$, in general, different in different relations.

2. Auxiliary assertions

Lemma 1. *Suppose* $S(t)$ *is a smooth complex-valued function for* $t_0 \leqslant t \leqslant t_k$, $t_0 < t_1 < \ldots < t_k$ *and* $\min\limits_{0 \leqslant j \leqslant k-1} (t_{j+1} - t_j) = \delta > 0$. *Then the following inequality holds*:

$$\sum_{j=1}^{k} |S(t_j)|^2 \leqslant \frac{1}{\delta} j_1 + 2\sqrt{j_1 j_2},$$

where

$$j_1 = \int_{t_0}^{t_k} |S(t)|^2 dt, \quad j_2 = \int_{t_0}^{t_k} |S'(t)|^2 dt.$$

For the proof, see [1, Chapter VII, §1].

Lemma 2. *Suppose* M, N, Q *are integers. Then for any sequence of complex numbers* a_n *the following estimation holds*:

$$\sum_{q \leqslant Q} \sideset{}{^*}\sum_{\chi \bmod q} \Bigl| \sum_{n=M+1}^{M+N} a_n \chi(n) \Bigr|^2 \ll (Q^2 + N) \sum_{n=M+1}^{M+N} |a_n|^2,$$

where the constant in the symbol $\ll$ *is absolute.*

For the proof, see [1, Chapter IX, answers to problems].

Lemma 3. *Suppose* $q \geqslant 3$, χ *is non-principle Dirichlet's character modulo* q. *Then for any* s, Y *such that* $Res \geqslant \sigma_0 > 0$ *and* $Y \geqslant q(|t|+1)/\pi$ *the following equality holds*:

$$L(s, \chi) = \sum_{n \leqslant Y} \frac{\chi(n)}{n^s} + O\bigl(qY^{-\sigma}\bigr),$$

where the constant in the symbol O *is absolute.*

For the proof, see [2, §26].

Lemma 4. *Let c be a sufficiently small positive absolute constant and let $q \geqslant 3$. Suppose χ is a complex character modulo q. Then the function $L(s,\chi)$ has no zeros in the domain*

$$\operatorname{Re} s > 1 - \frac{c}{\ln q(|t|+1)}, \quad -\infty < t < +\infty;$$

now if χ is a real non-principal Diriclet's character modulo q then the function $L(s,\chi)$ has no zeros in the domain

$$\operatorname{Re} s > 1 - \frac{c}{\ln q(|t|+1)}, \quad |t| > 0.$$

For the proof, see [1, Chapter IX, §2].

Lemma 5 (Siegel). *For any ε, $0 < \varepsilon < \frac{1}{2}$, there exists $c = c(\varepsilon) > 0$ such that if χ is a real character modulo q and β is a real zero of $L(s,\chi)$, then*

$$\beta < 1 - \frac{c}{q^{\varepsilon}}.$$

For the proof, see [1, Chapter IX, §2]. The constant $c = c(\varepsilon)$ is not effective. This means that it is impossible to find or estimate $c(\varepsilon)$ from a given ε. Therefore all statements (including main theorem of this paper) in which this lemma is essentially used are ineffective, too.

Lemma 6 (Montgomery). *For any $Q \geqslant 3$, $T \geqslant 3$ the following estimation holds:*

$$\sum_{q \leqslant Q} \sum_{\chi \bmod q}{}^{*} N(\sigma; T, \chi) \ll \bigl(Q^2 T\bigr)^{\vartheta(\sigma)} (\ln QT)^{14}$$

where

$$\vartheta(\sigma) = \begin{cases} \dfrac{3(1-\sigma)}{2-\sigma}, & \text{if } \ \frac{1}{2} \leqslant \sigma \leqslant \frac{4}{5}, \\ \dfrac{2(1-\sigma)}{\sigma}, & \text{if } \ \frac{4}{5} \leqslant \sigma \leqslant 1, \end{cases}$$

and the constant in the symbol $\ll$ is absolute.

For the proof, see [3, Chapter 12].

Lemma 7. *Suppose $q \geqslant 1$ is an integer, χ is a character modulo q and $\varrho = \beta + i\gamma$ runs through all non-trivial zeros of $L(s,\chi)$. Then*

$$\sum_{|\gamma| \leqslant T} \frac{1}{|\gamma| + 1} \ll (\ln qT)^2,$$

as $T \to +\infty$.

This estimation follows from asymptotic formula for $N(T,\chi)$ - the number of zeros of $L(s,\chi)$ in the rectangle $0 \leqslant \operatorname{Re} s \leqslant 1$, $|\operatorname{Im} s| \leqslant T$.

3. Basic assertions

Lemma 8. *Let χ be a non-primitive character modulo $q = q_1 r$ induced by primitive character χ_1 modulo q_1. Then*

$$|L(s,\chi)| \leqslant \tau(r)|L(s,\chi_1)|$$

in the half-plane $\mathrm{Re}\, s \geqslant 0$.

Proof. Using the formula

$$L(s,\chi) \;=\; L(s,\chi_1)\prod_{p|q,\, p\nmid q_1}\Bigl(1-\frac{\chi_1(p)}{p^s}\Bigr),$$

we get

$$|L(s,\chi)| \leqslant |L(s,\chi_1)|\prod_{p|r}\Bigl(1+\frac{1}{p^{\sigma}}\Bigr) \leqslant |L(s,\chi_1)|\prod_{p|r}2 \leqslant \tau(r)|L(s,\chi_1)|.$$

Lemma 9. *Suppose $f(n)$ is non-negative multiplicative function such that $f(n) = O(n^{\varepsilon})$, $0<\varepsilon<\frac{1}{2}$, and the function*

$$F(s) = \sum_{n=1}^{+\infty}\frac{f(n)}{n^s}$$

satisfies the identity

$$F(s) \;=\; \sqrt{\zeta(s)}\,\Phi(s)$$

for $\mathrm{Re}\, s > 1$ ($\sqrt{z} > 0$ *for* $z > 0$), *where $\Phi(s)$ is regular in the half-plane* $\mathrm{Re}\, s > \frac{1}{2}$ *and obeys the estimate*

$$|\Phi(\sigma+it)| \ll \max\Bigl\{1, \bigl(\sigma-\tfrac{1}{2}\bigr)^{-c}\Bigr\}$$

for any $\sigma > \frac{1}{2}$ and some $c > 0$. Then

$$\sum_{n\leqslant x} f(n) \;=\; \frac{x}{\sqrt{\ln x}}\biggl(\frac{\Phi(1)}{\sqrt{\pi}} \;+\; O\Bigl(\frac{1}{\ln x}\Bigr)\biggr)$$

where the constant in the symbol O depends on f only.

Proof. Since $f(n) = O(n^{\varepsilon})$, without loss of generality we may assume that x has the form $N+\frac{1}{2}$ for some integer N. Suppose T differs from the imaginary part of any zero of $\zeta(s)$ and $2\leqslant T\leqslant x$. Then, by Perron's formula we get

$$\sum_{n\leqslant x} f(n) \;=\; I \;+\; O\Bigl(\frac{x}{T}\ln x\Bigr)$$

where

$$I \;=\; \frac{1}{2\pi i}\int_{b-iT}^{b+iT} F(s)\,\frac{x^s}{s}\,ds, \quad b = 1+\frac{1}{\ln x}.$$

Let $c_1 > 0$ be small positive constant such that $\zeta(s)$ has no zeros in the rectangle with vertices $1 \pm iT$, $\alpha \pm iT$,

$$\alpha \;=\; 1 \;-\; c_1(\ln T)^{-\frac{2}{3}}(\ln\ln T)^{-\frac{1}{3}}.$$

Then, by the identity $F(s) = \sqrt{\zeta(s)}\,\Phi(s)$ the function $F(s)$ continues analytically to the domain $\alpha \leqslant \operatorname{Re} s \leqslant 1$, $|\operatorname{Im} s| \leqslant T$ with horizontal cut going straight from the point $s = \alpha$ to the point $s = 1$. By Cauchy's theorem,

$$I \;=\; -\sum_{k=1}^{6} I_k,$$

where the symbols $I_1, \ldots, I_4$ denote the integrals along the segments connecting points $b+iT$, $\alpha+iT$, α, $\alpha-iT$, $b-iT$ and the symbols I_5, I_6 denote the integrals along the upper and lower edges of the cut respectively.

Since the bound

$$\zeta(\sigma + it) \;=\; O\bigl(\ln^{\frac{2}{3}}(|t| + 2)\bigr)$$

holds along the contour, we obtain:

$$|I_1| + |I_4| \;\ll\; \int_{\alpha}^{b}\bigl(|F(\sigma + iT)| \;+\; |F(\sigma - iT)|\bigr)\frac{x^{\sigma}d\sigma}{\sqrt{\sigma^2 + T^2}} \;\ll\; \frac{x}{T}(\ln T)^{\frac{1}{3}},$$

$$|I_2| + |I_3| \ll \int_{-T}^{T}|F(\alpha + it)|\frac{x^{\alpha}\,dt}{\sqrt{\alpha^2 + t^2}} \ll x^{\alpha}\int_{0}^{T}\frac{\ln^{\frac{1}{3}}(t + 2)}{t + 2}dt \ll x^{\alpha}(\ln x)^{\frac{4}{3}}.$$

Consider the function $u(s) = (s-1)\zeta(s)$, $u(1) = 1$. Since $u(s) \neq 0$ for $|s-1| \leqslant \frac{1}{2}$, it follows that

$$\begin{aligned}
I_5 + I_6 &= \frac{1}{2\pi i}\int_{\alpha}^{1}\bigl(\sqrt{\zeta(\sigma + i\cdot 0)} \;-\; \sqrt{\zeta(\sigma - i\cdot 0)}\bigr)\Phi(\sigma)\,\frac{x^{\sigma}}{\sigma}d\sigma \;= \\
&= \frac{1}{2\pi i}\int_{\alpha}^{1}\Bigl(\frac{1}{\sqrt{\sigma - 1 + i\cdot 0}} - \frac{1}{\sqrt{\sigma - 1 - i\cdot 0}}\Bigr)\Phi(\sigma)\sqrt{u(\sigma)}\,\frac{x^{\sigma}}{\sigma}d\sigma \;= \\
&= \frac{1}{2\pi i}\int_{0}^{1-\alpha}\Bigl(\frac{1}{i\sqrt{v}} \;-\; \frac{1}{(-i\sqrt{v})}\Bigr)\,\frac{\Phi(1 - v)\sqrt{u(1 - v)}}{1 - v}\,x^{1-v}\,dv \;= \\
&= -\,\frac{x}{\pi}\int_{0}^{1-\alpha}\frac{x^{-v}}{\sqrt{v}}\,\frac{\Phi(1 - v)\sqrt{u(1 - v)}}{1 - v}\,dv.
\end{aligned}$$

The functions $\Phi(s)$ and $u(s)$ are bounded for $|s - 1| \leqslant \frac{1}{3}$. This implies that

$$\frac{\Phi(1 - v)\sqrt{u(1 - v)}}{1 - v} \;=\; \Phi(1) \;+\; O(|v|)$$

where the constant in the O-symbol depends on Φ only. Thus we obtain

$$\begin{aligned}
I_5 + I_6 &= -\,\frac{x}{\pi}\int_{0}^{1-\alpha}\frac{x^{-v}}{\sqrt{v}}\bigl(\Phi(1) \;+\; O(v)\bigr)dv \;= \\
&= -\,\frac{x}{\pi}\,\Phi(1)\Bigl(\int_{0}^{+\infty}\frac{x^{-v}}{\sqrt{v}}\,dv \;-\; \int_{1-\alpha}^{+\infty}\frac{x^{-v}}{\sqrt{v}}\,dv\Bigr) \;+\; O\Bigl(x\int_{0}^{+\infty}\sqrt{v}x^{-v}\,dv\Bigr).
\end{aligned}$$

Since

$$(1-\alpha)\ln x = c_1(\ln x)(\ln T)^{-\frac{2}{3}}(\ln\ln T)^{-\frac{1}{3}} \geqslant c_1\Bigl(\frac{\ln x}{\ln\ln x}\Bigr)^{\frac{1}{3}} > 1,$$

we get

$$\int_{1-\alpha}^{+\infty}\frac{x^{-v}}{\sqrt{v}}\,dv \;=\; \frac{1}{\sqrt{\ln x}}\int_{(1-\alpha)\ln x}^{+\infty}\frac{e^{-w}}{\sqrt{w}}\,dw < \frac{1}{\sqrt{\ln x}}\int_{(1-\alpha)\ln x}^{+\infty}e^{-w}\,dw = \frac{x^{\alpha-1}}{\sqrt{\ln x}},$$

$$I_5 + I_6 = -\,\frac{\Phi(1)}{\sqrt{\pi}}\,\frac{x}{\sqrt{\ln x}} \;+\; O\Bigl(\frac{x^{\alpha-1}}{\sqrt{\ln x}}\Bigr) + O\Bigl(\frac{x}{(\ln x)^{3/2}}\Bigr).$$

Using the above inequalities for $I_1,\ldots,I_4$ and substituting $T = e^{\sqrt{\ln x}}$, we finally obtain

$$I \;=\; \frac{\Phi(1)}{\sqrt{\pi}}\,\frac{x}{\sqrt{\ln x}}\Bigl(1 \,+\, O\Bigl(\frac{1}{\ln x}\Bigr)\Bigr).$$

The proof is complete.

Lemma 10. *Let m be an integer, $m \geqslant 1$. Then*

$$\sum_{\substack{q\leqslant x\\(q,m)=1}}\frac{1}{\varphi(q)} \;=\; C\beta(m)\Bigl(\ln x + \gamma - \sum_{p}\frac{\ln p}{p^2-p+1} + \sum_{p\mid m}\frac{p^2\ln p}{(p-1)(p^2-p+1)}\Bigr) +$$

$$+\,O\Bigl(\frac{\ln^2 x}{x}\Bigr) \;+\; O\Bigl(\frac{\tau(m)\ln x}{x}\Bigr),$$

where γ is Euler's constant,

$$C \;=\; \prod_{p}\Bigl(1 \,+\, \frac{1}{p(p-1)}\Bigr) \;=\; \frac{\zeta(2)\zeta(3)}{\zeta(6)},$$

and the constants in O's are absolute.

Proof. Note that

$$\frac{1}{\varphi(q)} = \frac{1}{q}\prod_{p\mid q}\Bigl(1-\frac{1}{p}\Bigr)^{-1} = \frac{1}{q}\prod_{p\mid q}\Bigl(1+\frac{1}{p-1}\Bigr) = \frac{1}{q}\prod_{p\mid q}\Bigl(1+\frac{1}{\varphi(p)}\Bigr) = \frac{1}{q}\sum_{d\mid q}\frac{\mu^2(d)}{\varphi(d)}.$$

Let us use the prime sign for the summation over the numbers coprime to m. Thus we get

$$\sum_{q\leqslant x}{}'\,\frac{1}{\varphi(q)} = \sum_{q\leqslant x}{}'\,\frac{1}{q}\sum_{d\mid q}\frac{\mu^2(d)}{\varphi(d)} = \sum_{d\leqslant x}{}'\,\frac{\mu^2(d)}{\varphi(d)}\sum_{\substack{q\leqslant x\\ q\equiv 0\,(\mathrm{mod}\,d)}}{}'\,\frac{1}{q} =$$

$$= \sum_{d\leqslant x}{}'\,\frac{\mu^2(d)}{\varphi(d)}\sum_{k\leqslant x/d}{}'\,\frac{1}{kd} = \sum_{d\leqslant x}{}'\,\frac{\mu^2(d)}{d\,\varphi(d)}\sum_{k\leqslant x/d}{}'\,\frac{1}{k} =$$

$$= \sum_{d\leqslant x}{}'\,\frac{\mu^2(d)}{d\,\varphi(d)}\sum_{k\leqslant x/d}\Bigl(\sum_{\delta\mid(k,m)}\mu(\delta)\Bigr)\frac{1}{k} = \sum_{d\leqslant x}{}'\,\frac{\mu^2(d)}{d\,\varphi(d)}\sum_{\delta\mid m}\mu(\delta)\sum_{\substack{k\leqslant x/d\\ k\equiv 0\,(\mathrm{mod}\,\delta)}}\frac{1}{k} =$$

$$= \sum_{\delta\mid m}\mu(\delta)\sum_{d\leqslant x}{}'\,\frac{\mu^2(d)}{d\,\varphi(d)}\sum_{r\leqslant x/(d\delta)}\frac{1}{r\delta} =$$

$$= \sum_{\delta\mid m}\frac{\mu(\delta)}{\delta}\sum_{d\leqslant x}{}'\,\frac{\mu^2(d)}{d\,\varphi(d)}\Bigl(\ln\frac{x}{d\delta} \,+\, \gamma \,+\, O\Bigl(\frac{d\delta}{x}\Bigr)\Bigr).$$

The contribution of the error term if the brackets to the initial sum does not exceed in order

$$\sum_{\delta|m}\frac{1}{\delta}\sideset{}{'}\sum_{d\leqslant x}\frac{1}{d\,\varphi(d)}\,\frac{d\delta}{x} \;=\; \frac{1}{x}\Bigl(\sum_{\delta|m}1\Bigr)\sum_{d\leqslant x}\frac{1}{\varphi(d)} \;=\; O\Bigl(\frac{\tau(m)\ln x}{x}\Bigr).$$

The contribution of all other terms has the form

$$\begin{aligned}
&\sum_{\delta|m}\frac{\mu(\delta)}{\delta}\sideset{}{'}\sum_{d\leqslant x}\frac{\mu^2(d)}{d\,\varphi(d)}\,(\ln x+\gamma-\ln d-\ln\delta) \;=\\
&= (\ln x+\gamma)\Bigl(\sum_{\delta|m}\frac{\mu(\delta)}{\delta}\Bigr)\sideset{}{'}\sum_{d\leqslant x}\frac{\mu^2(d)}{d\,\varphi(d)} \;-\; \Bigl(\sum_{\delta|m}\frac{\mu(\delta)}{\delta}\Bigr)\sideset{}{'}\sum_{d\leqslant x}\frac{\mu^2(d)\ln d}{d\,\varphi(d)} \;-\\
&- \Bigl(\sum_{\delta|m}\frac{\mu(\delta)\ln\delta}{\delta}\Bigr)\sideset{}{'}\sum_{d\leqslant x}\frac{\mu^2(d)}{d\,\varphi(d)}.
\end{aligned}$$

Let us replace all the sums over $d\leqslant x$ by infinite sums over d coprime to m. Using the inequality

$$\sum_{q\leqslant x}\frac{1}{\varphi(q)} \;\ll\; \ln x,$$

we obtain:

$$\begin{aligned}
\sideset{}{'}\sum_{d\leqslant x}\frac{\mu^2(d)}{d\,\varphi(d)} \;&=\; \sideset{}{'}\sum_{d=1}^{+\infty}\frac{\mu^2(d)}{d\,\varphi(d)} \;+\; O\Bigl(\frac{\ln x}{x}\Bigr),\\
\sideset{}{'}\sum_{d\leqslant x}\frac{\mu^2(d)\ln d}{d\,\varphi(d)} \;&=\; \sideset{}{'}\sum_{d=1}^{+\infty}\frac{\mu^2(d)\ln d}{d\,\varphi(d)} \;+\; O\Bigl(\frac{\ln^2 x}{x}\Bigr).
\end{aligned}$$

Since

$$\begin{aligned}
&\sideset{}{'}\sum_{d=1}^{+\infty}\frac{\mu^2(d)\ln d}{d\,\varphi(d)} \;=\; \prod_{p\nmid m}\Bigl(1\;+\;\frac{1}{p(p-1)}\Bigr) \;=\; \prod_{p}\Bigl(1\;+\;\frac{1}{p(p-1)}\Bigr)\prod_{p|m}\frac{p(p-1)}{p^2-p+1} \;=\\
&= C\prod_{p|m}\frac{p(p-1)}{p^2-p+1},
\end{aligned}$$

we have

$$\begin{aligned}
&\sideset{}{'}\sum_{n\leqslant x}\frac{1}{\varphi(q)} = C\,\frac{\varphi(m)}{m}\,(\ln x+\gamma)\prod_{p|m}\frac{p(p-1)}{p^2-p+1} \;-\; \frac{\varphi(m)}{m}\sideset{}{'}\sum_{d=1}^{+\infty}\frac{\mu^2(d)\ln d}{d\,\varphi(d)} \;-\\
&-\; C\prod_{p|m}\frac{p(p-1)}{p^2-p+1}\Bigl(\sum_{\delta|m}\frac{\mu(\delta)\ln\delta}{\delta}\Bigr).
\end{aligned}$$

Note that non-zero terms in the sum over $\delta|m$ correspond to squarefree divisors δ, so we have

$$\ln\delta \;=\; \sum_{p|\delta}\ln p.$$

Therefore,

$$\sum_{\delta|m} \frac{\mu(\delta)\ln\delta}{\delta} = \sum_{\delta|m} \frac{\mu(\delta)}{\delta} \sum_{p|\delta} \ln p = \sum_{p|m} \ln p \sum_{\substack{\delta|m \\ \delta\equiv 0 (\operatorname{mod} p)}} \frac{\mu(\delta)}{\delta} =$$

$$= \sum_{p|\delta} \ln p \sum_{\delta_1|\frac{m}{p},\, (\delta_1,p)=1} \frac{\mu(p)\mu(\delta_1)}{p\delta_1} = -\sum_{p|m} \frac{\ln p}{p} \sum_{\delta_1|\frac{m}{p},\, (\delta_1,p)=1} \frac{\mu(\delta_1)}{\delta_1} =$$

$$= -\sum_{p|m} \frac{\ln p}{p} \prod_{r|\frac{m}{p},\, r\neq p} \Big(1 - \frac{1}{r}\Big) = -\sum_{p|m} \frac{\ln p}{p} \cdot \frac{\displaystyle\prod_{p|m}\Big(1 - \frac{1}{r}\Big)}{1 - \dfrac{1}{p}} =$$

$$= -\frac{\varphi(m)}{m} \sum_{p|m} \frac{\ln p}{p-1}.$$

Next, by the same arguments we obtain

$$\sum_{d=1}^{+\infty}{}' \frac{\mu^2(d)\ln d}{d\,\varphi(d)} = \sum_{d=1}^{+\infty}{}' \frac{\mu^2(d)}{d\,\varphi(d)} \sum_{p|d} \ln p = \sum_{p}{}' \ln p \sum_{\substack{d=1 \\ d\equiv 0 (\operatorname{mod} p)}}^{+\infty}{}' \frac{\mu^2(d)}{d\,\varphi(d)} =$$

$$= \sum_{p}{}' \ln p \sum_{\substack{d_1=1 \\ (d_1,p)=1}}^{+\infty}{}' \frac{\mu^2(pd_1)}{pd_1\,\varphi(pd_1)} = \sum_{p}{}' \frac{\ln p}{p\,\varphi(p)} \sum_{\substack{d_1=1 \\ (d_1,pm)=1}}^{+\infty} \frac{\mu^2(d_1)}{d_1\,\varphi(d_1)} =$$

$$= \sum_{p}{}' \frac{\ln p}{p(p-1)} \prod_{q\nmid pm} \Big(1 + \frac{1}{q(q-1)}\Big) =$$

$$= \sum_{p}{}' \frac{\ln p}{p(p-1)} \cdot \frac{\displaystyle\prod_{q}\Big(1 + \frac{1}{q(q-1)}\Big)}{\Big(1 + \dfrac{1}{p(p-1)}\Big) \displaystyle\prod_{q|m}\Big(1 + \frac{1}{q(q-1)}\Big)} =$$

$$= C \sum_{p}{}' \frac{\ln p}{p(p-1)} \cdot \frac{p(p-1)}{p^2-p+1} \prod_{q|m} \Big(1 + \frac{1}{q(q-1)}\Big)^{-1} =$$

$$= C \sum_{p}{}' \frac{\ln p}{p^2-p+1} \prod_{q|m} \frac{q(q-1)}{q^2-q+1} =$$

$$= C \prod_{q|m} \frac{q(q-1)}{q^2-q+1} \Bigg(\sum_{p} \frac{\ln p}{p^2-p+1} - \sum_{p|m} \frac{\ln p}{p^2-p+1} \Bigg).$$

Substituting these relations in the above formula for the initial sum and taking into account the relation

$$\frac{\varphi(m)}{m} \prod_{p|m} \frac{p(p-1)}{p^2-p+1} = \prod_{p|m} \frac{(p-1)^2}{p^2-p+1} = \beta(m),$$

we get

$$\sum_{q\leqslant x}{}' \frac{1}{\varphi(q)} =$$

$$= C\beta(m)\Bigl(\ln x + \gamma - \sum_{p}\frac{\ln p}{p^2-p+1} + \sum_{p|a}\frac{\ln p}{p^2-p+1} + \sum_{p|a}\frac{\ln p}{p-1}\Bigr) +$$

$$+ O\Bigl(\frac{\ln^2 x}{x}\Bigr) + O\Bigl(\frac{\tau(a)\ln x}{x}\Bigr).$$

Corollary. *Under the same conditions,*

$$\sum_{q\leqslant x}{}' \frac{1}{\varphi(q)} = C\beta(m)\ln x + O\bigl(\tau(m)\bigr)$$

where the constant in the symbol O is absolute.

Lemma 11. *Let $d\geqslant 2$ be a fixed integer. Suppose δ runs over an increasing sequence that contains 1 and all integers which are not divisible by prime numbers coprime to d. Suppose also*

$$D_1(x) = \sum_{\delta\leqslant x} 1, \quad D_2(x) = \sum_{\delta > x}\frac{1}{\delta}.$$

Then

$$D_1(x) \ll (\ln x)^s \quad D_2(s) \ll \frac{(\ln x)^s}{x},$$

where $1\leqslant s\leqslant \tau(d)$.

Proof. Let $p_1^{\alpha_1}\dots p_s^{\alpha_s}$ be the unique decomposition of d into prime-powers. Then $D_1(x)$ equals to the number of solutions of the inequality $p^{\beta_1}\dots p_s^{\beta_s}\leqslant x$ or

$$\beta_1\ln p_1 + \ldots + \beta_s\ln p_s \leqslant \ln x$$

with non-negative integers $\beta_1,\dots,\beta_s$. Since $\ln p_1\geqslant\ln 2, \ldots, \ln p_s\geqslant \ln 2$, it follows that $D_1(x)$ does not exceed the number of solutions of the inequality

$$\beta_1 + \ldots + \beta_s \leqslant m, \quad m = \Bigl[\frac{\ln x}{\ln 2}\Bigr],$$

that is $D_1(x)\leqslant\binom{m+s}{s}$. Applying Stirling's formula and Cauchy's inequality we obtain

$$\binom{m+s}{s} = \frac{1}{s!}(m+s)\dots(m+1) \leqslant \frac{(m+s)^s}{s!} \leqslant \frac{2^{s-1}(m^s+s^s)}{(s/e)^s} =$$
$$= \frac{1}{2}\Bigl(\Bigl(\frac{2em}{s}\Bigr)^s + (2e)^s\Bigr) \leqslant (2em)^s \leqslant \Bigl(\frac{2e\ln x}{\ln 2}\Bigr)^s < (8\ln x)^s.$$

Further,

$$D_2(x) = \sum_{k=0}^{+\infty}\sum_{2^k x<\delta\leqslant 2^{k+1}x}\frac{1}{\delta} \leqslant \sum_{k=0}^{+\infty}\frac{1}{2^k x}\sum_{\delta\leqslant 2^{k+1}x} 1 \leqslant \sum_{k=0}^{+\infty}\frac{D_1(2^{k+1}x)}{2^k x} <$$
$$< \frac{1}{x}\sum_{k=0}^{+\infty}\frac{(8\ln(2^{k+1}x))^s}{2^k} \ll \frac{1}{x}\sum_{k=0}^{+\infty}\frac{(\ln x)^s + (k+1)^s}{2^k} \ll_s \frac{(\ln x)^s}{x}.$$

It remains to note that $s \leqslant \tau(d)$.

Lemma 12. *For any fixed* $a \geqslant 1$

$$S_a(x) \;=\; C(\ln x)\beta(a)E_a(x) \;+\; \theta R_a(x) \;+\; O(x\ln\ln x),$$

where the constant C *is defined in lemma 10,*

$$E_a(x) \;=\; \sum_{n\leqslant x}\frac{e_a(x)}{\tau(n)}, \quad e_a(n) \;=\; \frac{1}{\beta(a)}\sum_{d|(a,n)}\beta\Big(\frac{an}{d^2}\Big),$$

$$R_a(x) \;=\; \sum_{d|a}\;\sum_{\delta\leqslant(\ln x)^3} R_{a,d,\delta}(x),$$

$$R_{a,d,\delta}(x) \;=\; \sum_{\substack{q\leqslant y/d\\ (q,\frac{a\delta}{d})=1}}\frac{1}{\varphi(q)}\sum_{\chi\neq\chi_0}\Big|\sum_{\substack{m\leqslant\frac{x}{d\delta}\\ (m,d)=1}}\frac{\chi(m)}{\tau(m)}\Big|,$$

$y \;=\; \sqrt{x}(\ln x)^{-A}$, $A \;>\; 0$ *is arbitrary fixed number and* δ *runs through the sequence defined in lemma 11.*

Proof. By definition of $\tau(n)$,

$$S_a(x) \;=\; \sum_{uv\leqslant x}\frac{1}{\tau(uv+a)} \;=\; \Big(\sum_{u\leqslant\sqrt{x}}\;\sum_{v\leqslant\frac{x}{u}} \;+\; \sum_{\sqrt{x}<u\leqslant x}\;\sum_{v\leqslant\frac{x}{u}}\Big)\frac{1}{\tau(uv+a)} \;=$$

$$=\; \Big(\sum_{u\leqslant\sqrt{x}}\sum_{v\leqslant\frac{x}{u}} \;+\; \sum_{\sqrt{x}<u\leqslant x}\;\sum_{v\leqslant\frac{x}{u}}\Big)\frac{1}{\tau(uv+a)} \;=$$

$$=\; \Big(\sum_{u\leqslant\sqrt{x}}\sum_{v\leqslant\frac{x}{u}} \;+\; \sum_{v\leqslant\sqrt{x}}\sum_{u\leqslant\frac{x}{v}} \;-\; \sum_{u\leqslant\sqrt{x}}\sum_{v\leqslant\sqrt{x}}\Big)\frac{1}{\tau(uv+a)} \;=$$

$$=\; 2\sum_{u\leqslant\sqrt{x}}\sum_{v\leqslant\frac{x}{u}}\frac{1}{\tau(uv+a)} \;-\; \theta\,\tfrac{1}{2}\,x \;=\; 2\sum_{q\leqslant\sqrt{x}}\;\sum_{\substack{a<n\leqslant x+a\\ n\equiv a(\mathrm{mod}\,q)}}\frac{1}{\tau(n)} \;-\; \theta\,\tfrac{1}{2}\,x.$$

Replacing the domain of n in the inner sum to the interval $1\leqslant n\leqslant x$, we obtain

$$S_a(x) \;=\; 2\sum_{q\leqslant\sqrt{x}}\Big(\sum_{\substack{1\leqslant n\leqslant x\\ n\equiv a(\mathrm{mod}\,q)}} \;-\; \sum_{\substack{1\leqslant n\leqslant a\\ n\equiv a(\mathrm{mod}\,q)}} \;+\; \sum_{\substack{x<n\leqslant x+a\\ n\equiv a(\mathrm{mod}\,q)}}\Big)\frac{1}{\tau(n)} \;-\; \theta\,\tfrac{1}{2}\,x \;=$$

$$=\; 2\sum_{q\leqslant\sqrt{x}}\;\sum_{\substack{1\leqslant n\leqslant x\\ n\equiv a(\mathrm{mod}\,q)}}\frac{1}{\tau(n)} \;+\; 2\theta_1\sum_{q\leqslant\sqrt{x}}\Big(\frac{a}{q}+1\Big) \;-\; \theta\,\tfrac{1}{2}\,x \;=$$

$$=\; 2\sum_{q\leqslant\sqrt{x}}\;\sum_{\substack{1\leqslant n\leqslant x\\ n\equiv a(\mathrm{mod}\,q)}}\frac{1}{\tau(n)} \;+\; \theta_2 x.$$

Suppose A is an arbitrary positive number and let $y = \sqrt{x}(\ln x)^{-A}$. Then the summands corresponding to the values $y < q \leqslant \sqrt{x}$ do not exceed

$$2\sum_{y<q\leqslant\sqrt{x}}\;\sum_{\substack{n\leqslant x\\ n\equiv a(\mathrm{mod}\,q)}}\frac{1}{2} \leqslant \sum_{y<q\leqslant\sqrt{x}}\Big(\frac{x}{q} \;+\; 1\Big) \;<$$

$$<\; x\big(\ln\sqrt{x} \;-\; \ln y \;+\; O(y^{-1})\big) \;+\; \sqrt{x} \;=\; A\ln\ln x \;+\; 2\sqrt{x}(\ln x)^{A}.$$

Thus we get

$$S_a(x) = V_a(x) + 2\theta A \ln\ln x, \quad V_a(x) = 2\sum_{q\leqslant y}\sum_{\substack{1\leqslant n\leqslant x\\ n\equiv a (\operatorname{mod} q)}} \frac{1}{\tau(n)}.$$

Let us transform the last sum. Suppose $d = (q, a)$. Then $q = dq_1$, $a = da_1$ where $(q_1, a_1) = 1$. If $n \equiv a(\operatorname{mod} q)$ then for some $k \geqslant 0$ we obtain

$$n = a + kq = d(a_1 + kq_1) = dm,$$

where $m \equiv a_1(\operatorname{mod} q_1)$ and $m \leqslant x_d = x/d$. Thus the inner sum takes the form

$$\sum_{\substack{m\leqslant x_d\\ m\equiv a_1 (\operatorname{mod} q_1)}} \frac{1}{\tau(dm)}.$$

Since all possible values of d are among the divisors of a, it follows that

$$V_a(x) = 2\sum_{d|a}\sum_{\substack{q\leqslant y\\ (q,a)=d}}\sum_{\substack{m\leqslant x_d\\ m\equiv a_1 (\operatorname{mod} q_1)}} \frac{1}{\tau(dm)} =$$

$$= 2\sum_{d|a}\sum_{\substack{q_1\leqslant y_d\\ (q_1,a_1)=1}}\sum_{\substack{n\leqslant x_d\\ n\equiv a_1 (\operatorname{mod} q_1)}} \frac{1}{\tau(dn)},$$

where $y_d = y/d$.

Suppose d is an arbitrary fixed divisor of a and δ runs through an increasing sequence that contains 1 and positive integers all whose prime divisors are among prime divisors of d. (In particular, if $d = p$ is prime then δ take values $1, p, p^2, p^3, \ldots$). Then for any integer n there exists a unique representation in the form δm where δ belongs to the above sequence and $(m, d) = (m, \delta) = 1$. Obviously for such $n = \delta m$ we have $\tau(dn) = \tau(d\delta m) = \tau(d\delta)\tau(m)$. Thus we obtain

$$V_a(x) = 2\sum_{d|a}\sum_{\delta} V_{a,d,\delta}(x)$$

where

$$V_{a,d,\delta}(x) = \sum_{\substack{q_1\leqslant y_d\\ (q_1,a_1)=1}}\sum_{\substack{m\leqslant \frac{x_d}{\delta},\, (m,d)=1\\ \delta m\equiv a_1 (\operatorname{mod} q_1)}} \frac{1}{\tau(d\delta m)}.$$

Suppose that $(q_1, a_1) = 1$ and the congruence $\delta m \equiv a_1(\operatorname{mod} q_1)$ holds. If both the numbers δ and q_1 have the same divisor $\delta' > 1$ then δ' divides a_1. Therefore $(q_1, a_1) \geqslant \delta' > 1$. This contradiction shows that $(q_1, \delta) = 1$. Thus the solutions of the above congruence have the form $m \equiv a_1\delta^*(\operatorname{mod} q_1)$ This implies that

$$V_{a,d,\delta}(x) = \sum_{\substack{q_1\leqslant y_d\\ (q_1,a_1)=(q_1,\delta)=1}}\sum_{\substack{m\leqslant \frac{x_d}{\delta},\, (m,d)=1\\ m\equiv a_1\delta^* (\operatorname{mod} q_1)}} \frac{1}{\tau(d\delta m)}.$$

Let $\delta_0 = (\ln x)^3$. All the sums $V_{a,d,\delta}(x)$ corresponding to $\delta > \delta_0$ are estimated trivially:

$$V_{a,d,\delta}(x) \leqslant \sum_{q_1 \leqslant y_d} \sum_{\substack{m \leqslant \frac{x_d}{\delta} \\ m \equiv a_1\delta^* \,(\mathrm{mod}\, q_1)}} 1 \leqslant \sum_{q_1 \leqslant y_d} \Bigl(\frac{x_d}{\delta q_1} + 1\Bigr) \ll \frac{x}{d\delta}\ln x + \frac{y}{d}$$

where the constants in the symbols $\ll$ are absolute. By lemma 11, the contribution of these terms to $V_a(x)$ do not exceed in order

$$\sum_{d|a}\sum_{\delta>\delta_0}\Bigl(\frac{x}{d\delta}\ln x + \frac{y}{d}\Bigr) \ll x\ln x \Bigl(\sum_{d|a}\frac{1}{d}\Bigr)\sum_{\delta>\delta_0}\frac{1}{\delta} + y\Bigl(\sum_{d|a}\frac{1}{d}\Bigr)\Bigl(\sum_{\delta_0<\delta\leqslant x} 1\Bigr) \ll$$
$$\ll x\ln x\,\frac{(\ln\delta_0)^{\nu}}{\delta_0} + y(\ln x)^{\nu} \ll \frac{x(\ln\ln x)^{\nu}}{(\ln x)^2} \ll \frac{x}{\ln x},$$

where $\nu = \tau(a)$ and the constants in the symbols $\ll$ depend only on a. Thus,

$$V_a(x) = 2\sum_{d|a}\sum_{\delta\leqslant\delta_0} V_{a,d,\delta}(x) + O\Bigl(\frac{x}{\ln x}\Bigr).$$

Next, for $1 \leqslant \delta \leqslant \delta_0$ we have

$$V_{a,d,\delta}(x) = \sum_{\substack{q_1\leqslant y_d \\ (q_1,a_1\delta)=1}} \sum_{\substack{m\leqslant\frac{x_d}{\delta} \\ (m,d)=1}} \Bigl(\frac{1}{\varphi(q_1)}\sum_{\chi \,\mathrm{mod}\, q_1} \overline{\chi}(a_1\delta^*)\chi(m)\Bigr)\frac{1}{\tau(d\delta m)} =$$
$$= \sum_{\substack{q_1\leqslant y_d \\ (q_1,a_1\delta)=1}} \frac{1}{\varphi(q_1)} \sum_{\chi \,\mathrm{mod}\, q_1} \chi(\delta)\overline{\chi}(a_1) \sum_{\substack{m\leqslant\frac{x_d}{\delta} \\ (m,d)=1}} \frac{\chi(m)}{\tau(d\delta m)} =$$
$$= \sum_{\substack{q_1\leqslant y_d \\ (q_1,a_1\delta)=1}} \frac{1}{\varphi(q_1)} \sum_{\substack{m\leqslant\frac{x_d}{\delta} \\ (m,d)=(m,q_1)=1}} \frac{1}{\tau(d\delta m)} +$$
$$+ \frac{1}{\tau(d\delta)} \sum_{\substack{q_1\leqslant y_d \\ (q_1,a_1\delta)=1}} \frac{1}{\varphi(q_1)} \sum_{\substack{\chi \,\mathrm{mod}\, q_1 \\ \chi\neq\chi_0}} \chi(\delta)\overline{\chi}(a_1) \sum_{\substack{m\leqslant\frac{x_d}{\delta} \\ (m,d)=1}} \frac{\chi(m)}{\tau(m)}.$$

Therefore,

$$V_a(x) = W_a(x) + 2\theta R_a(x) + O\Bigl(\frac{x}{\ln x}\Bigr)$$

where

$$W_a(x) = 2\sum_{d|a}\sum_{\delta\leqslant\delta_0} \sum_{\substack{q_1\leqslant y_d \\ (q_1,a_1\delta)=1}} \frac{1}{\varphi(q_1)} \sum_{\substack{m\leqslant\frac{x_d}{\delta} \\ (m,d)=(m,q_1)=1}} \frac{1}{\tau(d\delta m)},$$

$$R_a(x) = \sum_{d|a}\sum_{\delta\leqslant\delta_0} R_{a,d,\delta}(x),$$

$$R_{a,d,\delta}(x) = \sum_{q\leqslant y_d} \frac{1}{\varphi(q_1)} \sum_{\substack{\chi \,\mathrm{mod}\, q \\ \chi\neq\chi_0}} \Bigl| \sum_{\substack{m\leqslant\frac{x_d}{\delta} \\ (m,d)=1}} \frac{\chi(m)}{\tau(m)} \Bigr|.$$

Let us replace the domain of δ in $W_a(x)$ by the segment $1\leqslant\delta\leqslant x_d$. Additional summands do not exceed

$$2\sum_{d|a}\sum_{\delta_0<\delta\leqslant x_d}\sum_{q_1\leqslant y_d}\frac{1}{\varphi(q_1)}\sum_{m\leqslant\frac{x_d}{\delta}}\frac{1}{2}\ll\sum_{d|a}\sum_{\delta_0<\delta\leqslant x_d}\frac{x}{d\delta}\ln x\ll$$

$$\ll x\ln x\Bigl(\sum_{d|a}\frac{1}{d}\Bigr)\sum_{\delta>\delta_0}\frac{1}{\delta}\ll x\ln x\,\frac{(\ln\delta_0)^{\nu}}{\delta_0}\ll\frac{x}{\ln x}.$$

Thus,

$$W_a(x)\,=\,2\sum_{d|a}\sum_{\delta}\sum_{\substack{q_1\leqslant y_d\\(q_1,a_1\delta)=1}}\frac{1}{\varphi(q_1)}\sum_{\substack{m\leqslant\frac{x_d}{\delta}\\(m,d)=(m,q_1)=1}}\frac{1}{\tau(d\delta m)}\,+\,O\Bigl(\frac{x}{\ln x}\Bigr).$$

Now our purpose is to transform double sum over δ and $m\leqslant\frac{x_d}{\delta}$ into single inner sum. We put $n=\delta m$. Since for any n, $1\leqslant n\leqslant x_d$, there exists a unique representation of the form $n=\delta m$ where $(m,d)=1$ and δ belongs to the above sequence, the condition $(m,d)=1$ in the inner sum in $W_a(x)$ may be omitted. Next, the condition $(m,q_1)=1$ should be replaced by the condition $(q_1,n)=1$. Indeed, the equality $(q_1,a_1\delta)=1$ in $W_a(x)$ implies that the sum over δ contains the terms that obey the condition $(q_1,\delta)=1$. Then both the conditions $(q_1,m)=1$ and $(q_1,\delta)=1$ are equivalent to the single condition $(q_1,n)=1$. Thus we get

$$W_a(x)\,=\,2\sum_{d|a}\sum_{\substack{q_1\leqslant y_d\\(q_1,a_1)=1}}\frac{1}{\varphi(q_1)}\sum_{\substack{n\leqslant x_d\\(q_1,n)=1}}\frac{1}{\tau(dn)}\,+\,O\Bigl(\frac{x}{\ln x}\Bigr)\,=$$

$$=\,2\sum_{d|a}\sum_{n\leqslant x_d}\frac{1}{\tau(dn)}\sum_{\substack{q_1\leqslant y_d\\(q_1,na_1)=1}}\frac{1}{\varphi(q_1)}\,+\,O\Bigl(\frac{x}{\ln x}\Bigr).$$

Applying the consequence of Lemma 9, we obtain

$$W_a(x)\,=\,2\sum_{d|a}\sum_{n\leqslant x_d}\frac{1}{\tau(dn)}\bigl(C\beta(na_1)\ln y_d\,+\,O\bigl(\tau(na_1)\bigr)\bigr)\,+\,O\Bigl(\frac{x}{\ln x}\Bigr).$$

Since $C\beta(na_1)\ln d+O\bigl(\tau(na_1)\bigr)\ll_a\tau(n)$, it follows that

$$W_a(x)\,=$$

$$=\,2\sum_{d|a}\sum_{n\leqslant x_d}\frac{1}{\tau(dn)}\bigl(C\beta(na_1)\bigl(\tfrac{1}{2}\ln x\,-\,A\ln\ln x\bigr)\,+\,O\bigl(\tau(n)\bigr)\bigr)\,+\,O\Bigl(\frac{x}{\ln x}\Bigr)\,=$$

$$=\,C(\ln x)\sum_{d|a}\sum_{n\leqslant x_d}\frac{\beta(na_1)}{\tau(dn)}\,+\,r_a(x),$$

where

$$
\begin{aligned}
r_a(x) \,&\ll\, (\ln\ln x)\sum_{d|a}\sum_{n\leqslant x_d}\frac{1}{\tau(dn)} \,+\, \sum_{d|a}\sum_{n\leqslant x_d}\frac{\tau(n)}{\tau(dn)} \,+\, \frac{x}{\ln x} \,\ll \\
&\ll\, (\ln\ln x)\Bigl(\sum_{d|a}1\Bigr)\sum_{m\leqslant x}\frac{1}{\tau(m)} \,+\, \sum_{d|a}\frac{x}{d} \,+\, \frac{x}{\ln x} \,\ll \\
&\ll\, \frac{x\ln\ln x}{\sqrt{\ln x}} \,+\, x \,+\, \frac{x}{\ln x} \,\ll\, x.
\end{aligned}
$$

Thus,

$$
W_a(x) \,=\, C(\ln x)\sum_{d|a}\sum_{n\leqslant x/d}\frac{\beta\Bigl(\dfrac{na}{d}\Bigr)}{\tau(nd)} \,+\, O(x).
$$

Changing the order of summation, we deduce that

$$
\begin{aligned}
W_a(x) \,&=\, C(\ln x)\sum_{m\leqslant x}\frac{1}{\tau(m)}\sum_{d|a,\ d|m}\beta\Bigl(\frac{na}{d}\Bigr) \,+\, O(x) \,= \\
&=\, C(\ln x)\,\beta(a)\sum_{m\leqslant x}\frac{e_a(m)}{\tau(m)} \,+\, O(x),
\end{aligned}
$$

where

$$
e_a(m) \,=\, \frac{1}{\beta(a)}\sum_{d|(a,m)}\beta\Bigl(\frac{na}{d^2}\Bigr)
$$

This completes the proof of the lemma.

The following lemma is the main assertion of the paper.

Lemma 13. *Let d be a fixed integer, $d\geqslant 1$. Then for any $B>0$ there exists $A=A(B)>0$ such that the estimation*

$$
R \,=\, \sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\substack{\chi \bmod q\\ \chi\neq\chi_0}}\Bigl|\sum_{\substack{n\leqslant N\\ (n,d)=1}}\frac{\chi(n)}{\tau(n)}\Bigr| \,\ll\, x(\ln x)^{-B}
$$

holds for any Q, N such that $Q\leqslant\sqrt{x}(\ln x)^{-A}$, $N\leqslant x$.

Remark. This assertion holds true for $A=\frac{16}{3}B+4$.

Proof. First we prove that if T do not coincide with an ordinate of a zero of $L(s,\chi)$ then the following inequality holds:

$$
\begin{aligned}
&\Bigl|\sum_{\substack{n\leqslant N\\ (n,d)=1}}\frac{\chi(n)}{\tau(n)}\Bigr| \,\ll \\
&\ll\, (\ln x)^{\alpha}\biggl(\frac{N}{T}\ln x \,+\, \frac{N}{T}\int_{\sigma_1}^{\sigma_2}\bigl(|L(\sigma+iT,\chi)|^{\frac{1}{2}} \,+\, |L(\sigma+iT,\overline{\chi})|^{\frac{1}{2}}\bigr)d\sigma \,+ \\
&+\, \sqrt{N}\int_0^T\bigl(|L(\sigma_1+it,\chi)|^{\frac{1}{2}} \,+\, |L(\sigma_1+it,\overline{\chi})|^{\frac{1}{2}}\bigr)\frac{dt}{t+1} \,+ \\
&+\, \sum_{|\gamma|\leqslant T}\frac{1}{|\gamma|+1}\int_{\sigma_1}^{\beta}|L(\sigma+i\gamma,\chi)|^{\frac{1}{2}}N^{\sigma}d\sigma\biggr).
\end{aligned}
$$

Here $\alpha = \frac{1}{24}$, $\sigma_1 = \frac{1}{2} + (\ln x)^{-1}$, $\sigma_2 = \sigma_1 + \frac{1}{2}$ and $\varrho = \beta + i\gamma$ runs through all zeros of $L(s, \chi)$ in the rectangle $\sigma_1 < \beta \leqslant 1$, $|\gamma| \leqslant T$. Assume that $\operatorname{Re} s > 1$. Then the generating function $F(s; \chi, d) = F(s)$ of the sequence $\chi(n)/\tau(n)$, $(n, d) = 1$, has the form

$$F(s) \,=\, \sum_{\substack{n=1\\(n,d)=1}}^{+\infty} \frac{\chi(n)}{\tau(n)}\, n^{-s} \,=\, \prod_{p\nmid d} F_p(s), \quad F_p(s) \,=\, 1 \,+\, \sum_{k=1}^{+\infty} \frac{\chi(p^k)}{k+1}\, p^{-ks}.$$

Taking for brevity $\kappa = \chi(p)p^{-s}$ we obtain for $\sigma > 1$:

$$F_p(s) \,=\, 1 \,+\, \tfrac{1}{2}\kappa \,+\, \tfrac{1}{3}\kappa^2 \,+\, \tfrac{1}{4}\kappa^3 \,+ \ldots =\, (1-\kappa)^{-\frac{1}{2}}(1-\kappa^2)^{\frac{1}{24}}(1-\kappa^3)^{\frac{1}{24}} f_p(s)$$

where the symbol $f_p(s)$ denotes a convergent series of the form

$$1 \,-\, \tfrac{49}{2880}\kappa^4 \,-\, \tfrac{49}{1440}\kappa^5 \,+\, \tfrac{1447}{362880}\kappa^6 \,-\, \tfrac{3383}{120360}\kappa^7 \,+ \ldots .$$

Therefore,

$$\begin{aligned}
F(s) \,&=\, \prod_{p\nmid d}\Bigl(1 \,-\, \frac{\chi(p)}{p^s}\Bigr)^{-\frac{1}{2}}\Bigl(1 \,-\, \frac{\chi^2(p)}{p^{2s}}\Bigr)^{\frac{1}{24}}\Bigl(1 \,-\, \frac{\chi^3(p)}{p^{3s}}\Bigr)^{\frac{1}{24}} f_p(s) \,= \\
&=\, \bigl(L(s,\chi)\bigr)^{\frac{1}{2}}\bigl(L(2s,\chi^2)\bigr)^{-\frac{1}{24}}\bigl(L(3s,\chi^3)\bigr)^{-\frac{1}{24}} \times \\
&\times \prod_{p\mid d}\Bigl(1 \,-\, \frac{\chi(p)}{p^s}\Bigr)^{\frac{1}{2}}\Bigl(1 \,-\, \frac{\chi^2(p)}{p^{2s}}\Bigr)^{-\frac{1}{24}}\Bigl(1 \,-\, \frac{\chi^3(p)}{p^{3s}}\Bigr)^{-\frac{1}{24}} \prod_{p\nmid d} f_p(s) \,= \\
&=\, \frac{\bigl(L(s,\chi)\bigr)^{\frac{1}{2}}}{\bigl(L(2s,\chi^2)L(3s,\chi^3)\bigr)^{\alpha}}\,\Phi(s),
\end{aligned}$$

where $\alpha = \frac{1}{24}$ and the function $\Phi(s)$ is regular in the half-plane $\operatorname{Re} s > \frac{1}{4}$. Suppose $T \geqslant 2$ do not coincide with an ordinate of a zero of $L(s, \chi)$. Taking $N_1 = [N] + \frac{1}{2}$, $\sigma_2 = 1 + (\ln x)^{-1}$, by Perron's formula we get

$$\sum_{\substack{n\leqslant N\\(n,d)=1}} \frac{\chi(n)}{\tau(n)} \,=\, \frac{1}{2\pi i}\int_{\sigma_2 - iT}^{\sigma_2 + iT} F(s)\frac{N_1^s}{s}\,ds \,+\, O\Bigl(\frac{N}{T}\ln x\Bigr) \,+\, O(1).$$

Suppose $\varrho = \beta + i\gamma$ runs through all the zeros of $L(s, \chi)$ in the domain $\sigma_2 < \beta \leqslant 1$, $|\gamma| \leqslant T$. Let Γ be a boundary of the rectangle with vertices $\sigma_1 \pm iT$, $\sigma_2 \pm iT$ and with horizontal cuts going from the left side of the rectangle to each zero ϱ. Applying Cauchy's theorem we obtain

$$\frac{1}{2\pi i}\int_{\sigma_2 - iT}^{\sigma_2 + iT} F(s)\frac{N_1^s}{s}\,ds \,=\, -\Bigl(I_1 \,+\, I_2 \,+\, I_3 \,+ \sum_{|\gamma|\leqslant T} I(\varrho)\Bigr)$$

where the symbols I_1, I_2, I_3 denote the integrals

$$\frac{1}{2\pi i}\int_{\sigma_2 + iT}^{\sigma_1 + iT} F(s)\frac{N_1^s}{s}\,ds, \;\; \frac{1}{2\pi i}\int_{\sigma_1 - iT}^{\sigma_2 - iT} F(s)\frac{N_1^s}{s}\,ds, \;\; \frac{1}{2\pi i}\,\text{v.p.}\int_{\sigma_1 + iT}^{\sigma_1 - iT} F(s)\frac{N_1^s}{s}\,ds$$

respectively, and $I(\varrho)$ means the sum of the integrals over the upper and lower edges of the cut:

$$I(\varrho) \,=\, \frac{1}{2\pi i}\biggl(\int_{\sigma_1+i(\gamma+0)}^{\beta+i(\gamma+0)} \,+\, \int_{\beta+i(\gamma-0)}^{\sigma_1+i(\gamma-0)}\biggr)F(s)\frac{N_1^s}{s}\,ds.$$

Since the infinite product for $\Phi(s)$ converges absolutely in the half-plane $\operatorname{Re} s > \frac{1}{4}$, then $|\Phi(s)| = O(1)$ along the contour Γ. Moreover, for $\sigma \geqslant \sigma_1$ we obtain

$$\begin{aligned}
&|L(3s,\chi^3)|^{-1} \,=\, \Bigl|\sum_{n=1}^{+\infty}\frac{\mu(n)\chi^3(n)}{n^{3s}}\Bigr| \,\leqslant\, \sum_{n=1}^{+\infty} n^{-3/2} \,=\, O(1),\\
&|L(2s,\chi^2)|^{-1} \,=\, \Bigl|\sum_{n=1}^{+\infty}\frac{\mu(n)\chi^2(n)}{n^{2s}}\Bigr| \,\leqslant\, \sum_{n=1}^{+\infty} n^{-2\sigma} \,\leqslant\, \sum_{n=1}^{+\infty} n^{2\sigma_1} \,\leqslant\\
&\leqslant\, 1 \,+\, \int_1^{+\infty} u^{-2\sigma_1}du \,=\, \frac{2\sigma_1}{2\sigma_1-1} \,=\, \tfrac{1}{2}\ln x + 1 \,<\, \ln x.
\end{aligned}$$

Thus,

$$|F(s)| \,\ll\, (\ln x)^{\alpha}|L(s,\chi)|^{\frac{1}{2}}$$

along the contour Γ. Therefore,

$$\begin{aligned}
&I_1 \ll \int_{\sigma_1}^{\sigma_2} |F(\sigma+iT)|\frac{N_1^{\sigma}d\sigma}{\sqrt{\sigma^2+T^2}} \,\ll\, \frac{N^{\sigma_2}}{T}(\ln x)^{\alpha}\int_{\sigma_1}^{\sigma_2} |L(\sigma+iT,\chi)|^{\frac{1}{2}}d\sigma \,\ll\\
&\ll\, \frac{N}{T}(\ln x)^{\alpha}\int_{\sigma_1}^{\sigma_2} |L(\sigma+iT,\chi)|^{\frac{1}{2}}d\sigma,\\
&I_2 \,\ll\, \frac{N}{T}(\ln x)^{\alpha}\int_{\sigma_1}^{\sigma_2} |L(\sigma+iT,\overline{\chi})|^{\frac{1}{2}}d\sigma.
\end{aligned}$$

Next,

$$\begin{aligned}
&|I_3| \,=\, \frac{1}{2\pi}\Bigl|\text{v.p.}\int_{-T}^{T} F(\sigma_1+it)\,\frac{N_1^{\sigma_1+it}}{\sigma_1+it}\,dt\Bigr| \,\ll\, N^{\sigma_1}\int_{-T}^{T}\frac{|F(\sigma_1+it)|dt}{\sqrt{\sigma_1^2+t^2}} \,\ll\\
&\ll\, \sqrt{N}(\ln x)^{\alpha}\int_{-T}^{T}\frac{|L(\sigma_1+it,\chi)|^{\frac{1}{2}}dt}{\sqrt{\sigma_1^2+t^2}} \,\ll\\
&\ll\, \sqrt{N}(\ln x)^{\alpha}\int_0^T \bigl(L(\sigma_1+it,\chi)|^{\frac{1}{2}} \,+\, L(\sigma_1+it,\overline{\chi})|^{\frac{1}{2}}\bigr)\frac{dt}{t+1}.
\end{aligned}$$

Finally, each of the integrals $I(\varrho)$ obeys the inequality

$$|I(\varrho)| \,\ll\, \frac{(\ln x)^{\alpha+1}}{|\gamma|+1}\int_{\sigma_1}^{\beta} |L(\sigma_1+i\gamma,\chi)|^{\frac{1}{2}}N^{\sigma}d\sigma.$$

Summing the above estimates we obtain the required inequality for the sum of $\chi(n)/\tau(n)$, $n \leqslant N$, $(n,d) = 1$. Summing these inequalities over all non-principle characters $\chi \bmod q$ and over $q \leqslant Q$, we get

$$R \,\ll\, (\ln x)^{\alpha}\sum_{j=1}^{4} R_j,$$

where

$$\begin{aligned}
R_1 &= \sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\chi\ne\chi_0}\frac{N}{T}\ln x,\\
R_2 &= \sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\chi\ne\chi_0}\frac{N}{T}\int_{\sigma_1}^{\sigma_2}\bigl(|L(\sigma+iT,\chi)|^{\frac{1}{2}} + |L(\sigma+iT,\overline{\chi})|^{\frac{1}{2}}\bigr)d\sigma,\\
R_3 &= \sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\chi\ne\chi_0}\sqrt{N}\int_0^T\bigl(|L(\sigma_1+it,\chi)|^{\frac{1}{2}} + |L(\sigma_1+it,\overline{\chi})|^{\frac{1}{2}}\bigr)\frac{dt}{t+1},\\
R_4 &= \sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\chi\ne\chi_0}\sum_{|\gamma|\leqslant T}\frac{1}{|\gamma|+1}\int_{\sigma_1}^{\beta}N^{\sigma}|L(\sigma+i\gamma,\overline{\chi})|^{\frac{1}{2}}d\sigma.
\end{aligned}$$

The values of T depends on q and χ only: $T=T(q,\chi)$. We choose $T(q,\chi)$ in a following way. The points $Q\cdot 2^{-k}$, $k=0,1,2,\ldots$ split the segment $(1,Q]$ into the intervals of the type $(M,M_1]$, where $M_1\leqslant 2M$, $M_1\leqslant Q$. Suppose the character χ modulo q is induced by a primitive character χ_1 modulo q_1, $q=q_1r$, where $M<q_1\leqslant M_1$. Then we put $T=M(\ln x)^C$, where the constant C, $0<C<A$, will be chosen later. Therefore, it follows that $T(q,\chi)=T(q_1,\chi_1)$; in particular, $T(q,\chi)$ do not depends on r. Replacing (if it is necessary) the value $T(q,\chi)$ by the value $T(q,\chi)+h$ for some h, $0<h\leqslant c(\ln x)^{-1}$, we may assume that $T(q,\chi)$ does not coincide with an ordinate of a zero of $L(s,\chi)$. In each case, we obviously have

$$M(\ln x)^C\leqslant T(q,\chi)\leqslant M(\ln x)^C+c(\ln x)^{-1}\leqslant 2M(\ln x)^C.$$

In the following, the sums over $\chi \bmod q$, $\chi\ne\chi_0$ are replaced by the sums over primitive characters $\chi_1 \bmod q_1$.

1°. Estimation of R_1. Obviously we have

$$\begin{aligned}
R_1 &\ll N(\ln x)\sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\chi\ne\chi_0}\frac{1}{T(q,\chi)}\ll\\
&\ll N(\ln x)\sum_{q_1\leqslant Q}\sum_{r\leqslant\frac{Q}{q_1}}\frac{1}{\varphi(q_1r)}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\frac{1}{T(q_1,\chi_1)}.
\end{aligned}$$

Using the symbol $\sum'_{M\leqslant Q}$ for the summation over the points $M=Q\cdot 2^{-k}$, by the inequality $\varphi(q_1r)\geqslant\varphi(q_1)\,\varphi(r)$ we obtain

$$\begin{aligned}
R_1 &\ll N(\ln x)\sideset{}{'}\sum_{M\leqslant Q}\sum_{M<q_1\leqslant M_1}\frac{1}{\varphi(q_1)}\sum_{r\leqslant\frac{Q}{q_1}}\frac{1}{\varphi(r)}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\frac{1}{M(\ln x)^C}\ll\\
&\ll\frac{N\ln x}{(\ln x)^C}\sideset{}{'}\sum_{M\leqslant Q}\frac{1}{M}\sum_{M<q_1\leqslant M_1}\frac{1}{\varphi(q_1)}\Bigl(\sideset{}{^*}\sum_{\chi_1 \bmod q_1}1\Bigr)\sum_{r\leqslant Q}\frac{1}{\varphi(r)}\ll\\
&\ll\frac{N(\ln x)^2}{(\ln x)^C}\sideset{}{'}\sum_{M\leqslant Q}\frac{1}{M}\Bigl(\sum_{M<q_1\leqslant M_1}1\Bigr)\ll\frac{N(\ln x)^2}{(\ln x)^C}\sideset{}{'}\sum_{M\leqslant Q}1\ll\frac{N(\ln x)^3}{(\ln x)^C}.
\end{aligned}$$

2°. Estimation of R_2. Since $\overline{\chi}$ and χ run through the same set of characters, we get

$$R_2 \ll N\int_{\sigma_1}^{\sigma_2} r_2(\sigma)d\sigma,$$

where

$$r_2(\sigma) = \sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\substack{\chi \bmod q\\ \chi\neq\chi_0}}\frac{|L(\sigma+iT,\chi)|^{\frac{1}{2}}}{T(q,\chi)}.$$

Using the same arguments as above, by lemma 7 we obtain ($T = T(q_1,\chi_1)$):

$$r_2(\sigma) \leqslant \sum_{q_1\leqslant Q}\frac{1}{\varphi(q_1)}\sum_{r\leqslant\frac{Q}{q_1}}\frac{1}{\varphi(r)}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\frac{\sqrt{\tau(r)}}{T(q_1,\chi_1)}|L(\sigma+iT,\chi_1)|^{\frac{1}{2}} \leqslant$$

$$\leqslant \sum_{q_1\leqslant Q}\frac{1}{\varphi(q_1)}\sum_{r\leqslant\frac{Q}{q_1}}\frac{\sqrt{\tau(r)}}{\varphi(r)}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\frac{|L(\sigma+iT,\chi_1)|^{\frac{1}{2}}}{T(q_1,\chi_1)} \ll$$

$$\ll \sideset{}{'}\sum_{M\leqslant Q}\sum_{M<q_1\leqslant M_1}\frac{1}{\varphi(q_1)}\sum_{r\leqslant\frac{Q}{q_1}}\frac{\sqrt{\tau(r)}}{\varphi(r)}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\frac{|L(\sigma+iT,\chi_1)|^{\frac{1}{2}}}{M(\ln x)^C} \ll$$

$$\ll (\ln x)^{-C}\sideset{}{'}\sum_{M\leqslant Q}\frac{1}{M}\sum_{M<q_1\leqslant M_1}\frac{1}{\varphi(q_1)}\sum_{r\leqslant\frac{Q}{q_1}}\frac{\sqrt{\tau(r)}}{\varphi(r)}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}|L(\sigma+iT,\chi_1)|^{\frac{1}{2}}.$$

By the inequalities

$$\sum_{r\leqslant X}\sqrt{\tau(r)}\leqslant\Bigl(X\sum_{r\leqslant X}\tau(r)\Bigr)^{\frac{1}{2}} \ll X\sqrt{\ln X}, \quad \frac{1}{\varphi(r)} \ll \frac{\ln\ln(r+3)}{r},$$

we deduce that

$$\sum_{r\leqslant\frac{Q}{q_1}}\frac{\sqrt{\tau(r)}}{\varphi(r)} \ll (\ln\ln Q)\sum_{r\leqslant Q}\frac{\sqrt{\tau(r)}}{r} \ll$$

$$\ll (\ln\ln x)\Bigl(\frac{1}{Q}\sum_{r\leqslant Q}\sqrt{\tau(r)} + \int_2^Q\Bigl(\sum_{r\leqslant u}\sqrt{\tau(r)}\Bigr)\frac{du}{u^2}\Bigr) \ll (\ln\ln x)\int_2^Q\sqrt{\ln u}\,\frac{du}{u}$$

$$\ll (\ln x)^{\frac{3}{2}}\ln\ln x$$

and therefore

$$r_2(\sigma) \ll \frac{(\ln x)^{\frac{3}{2}}\ln\ln x}{(\ln x)^C}\sideset{}{'}\sum_{M\leqslant Q}\frac{1}{M}\sum_{M<q_1\leqslant M_1}\frac{1}{\varphi(q_1)}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}|L(\sigma+iT,\chi_1)|^{\frac{1}{2}} \ll$$

$$\ll \frac{(\ln x)^{\frac{3}{2}}(\ln\ln x)^2}{(\ln x)^C}\sideset{}{'}\sum_{M\leqslant Q}\frac{1}{M^2}\,r_2(\sigma,M),$$

where

$$r_2(\sigma, M) \;=\; \sum_{M<q_1\leqslant M_1}\;\sideset{}{^*}\sum_{\chi_1 \bmod q_1} |L(\sigma+iT,\chi_1)|^{\frac{1}{2}}.$$

Applying Hölder's inequality, we obtain

$$r_2^4(\sigma, M) \;\ll\; \Bigl(\sum_{M<q_1\leqslant M_1}\;\sideset{}{^*}\sum_{\chi_1 \bmod q_1} 1\Bigr)^3 \sum_{M<q_1\leqslant M_1}\;\sideset{}{^*}\sum_{\chi_1 \bmod q_1} |L(\sigma+iT,\chi_1)|^2 \;\ll$$
$$\ll\; M^6 \sum_{M<q_1\leqslant M_1}\;\sideset{}{^*}\sum_{\chi_1 \bmod q_1} |L(\sigma+iT,\chi_1)|^2.$$

Let $Y = 2M_1T(q_1,\chi_1)$; then for any q_1 such that $M < q_1 \leqslant M_1$, the following inequality holds: $Y \geqslant q_1(T(q_1,\chi_1)+1)/\pi$. By lemma 3, we get

$$|L(\sigma+iT,\chi_1)|^2 \;\ll\; \Bigl|\sum_{n\leqslant Y}\frac{\chi(n)}{n^{\sigma+iT}}\Bigr|^2 \;+\; (q_1Y^{-\sigma})^2.$$

Since

$$q_1Y^{-\sigma} \leqslant \frac{M_1}{\sqrt{Y}} \;=\; \frac{M_1}{\sqrt{2M_1T(q_1,\chi_1)}} \;<\; \sqrt{\frac{M_1}{T(q_1,\chi_1)}} \;<$$
$$<\; \sqrt{\frac{2M}{M(\ln x)^C}} \;=\; \sqrt{2}(\ln x)^{-C/2} \;<\; 1,$$

for $\sigma \geqslant \frac{1}{2}$, we get

$$r_2^4(\sigma, M) \;\ll\; M^8 \;+\; M^6 \sum_{q_1\leqslant M_1}\;\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\Bigl|\sum_{n\leqslant Y}\frac{\chi(n)}{n^{\sigma+iT}}\Bigr|^2.$$

Setting $Q = M_1$, $M = 0$, $N = Y$ in lemma 2, we obtain

$$r_2^4(\sigma, M) \;\ll\; M^8 \;+\; M^6(M^2 \;+\; Y)\sum_{n\leqslant Y}\frac{1}{n^{2\sigma}} \;\ll\; M^8(\ln x)^{C+1}$$

and therefore

$$r_2(\sigma) \;\ll\; \frac{(\ln x)^{\frac{3}{2}}(\ln\ln x)^2}{(\ln x)^C}\sideset{}{'}\sum_{M\leqslant Q}\frac{1}{M^2}\,M^2(\ln x)^{\frac{1}{4}(C+1)} \;\ll$$
$$\ll\; \frac{(\ln x)^{\frac{7}{4}}(\ln\ln x)^2}{(\ln x)^{\frac{3}{4}C}}\sideset{}{'}\sum_{M\leqslant Q} 1 \;\ll\; \frac{(\ln x)^{\frac{11}{4}}(\ln\ln x)^2}{(\ln x)^{\frac{3}{4}C}}.$$

Thus,

$$R_2 \;\ll\; N\,\frac{(\ln x)^{\frac{11}{4}}}{(\ln x)^{\frac{3}{4}C}}\,(\ln\ln x)^2.$$

3°. Estimation of R_3. By the same arguments we obtain

$$R_3 \ll \sqrt{N}\sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\substack{\chi \bmod q\\ \chi\neq\chi_0}}\int_0^{T(q,\chi)}|L(\sigma_1+it,\chi)|^{\frac{1}{2}}\,\frac{dt}{t+1} \ll$$

$$\ll \sqrt{N}\sum_{q_1\leqslant Q}\frac{1}{\varphi(q_1)}\sum_{r\leqslant \frac{Q}{q_1}}\frac{\sqrt{\tau(r)}}{\varphi(r)}\sum^{*}_{\chi_1 \bmod q_1}\int_0^{T(q_1,\chi_1)}|L(\sigma_1+it,\chi)|^{\frac{1}{2}}\,\frac{dt}{t+1}.$$

Since the terms in the inner sum over χ_1 do not depend on r, we easily get

$$R_3 \ll \sqrt{N}(\ln x)^{\frac{3}{2}}(\ln\ln x)\sum_{q_1\leqslant Q}\frac{1}{\varphi(q_1)}\sum^{*}_{\chi_1 \bmod q_1}\int_0^{T(q_1,\chi_1)}|L(\sigma_1+it,\chi)|^{\frac{1}{2}}\,\frac{dt}{t+1} \ll$$

$$\ll \sqrt{N}(\ln x)^{\frac{3}{2}}(\ln\ln x)\sum'_{M\leqslant Q}\sum_{M<q_1\leqslant M_1}\frac{1}{\varphi(q_1)}\sum^{*}_{\chi_1 \bmod q_1}\int_0^{T(q_1,\chi_1)}\frac{|L(\sigma_1+it,\chi)|^{\frac{1}{2}}dt}{t+1}$$

$$\ll \sqrt{N}(\ln x)^{\frac{3}{2}}(\ln\ln x)^2\sum'_{M\leqslant Q}\frac{1}{M}\int_0^{T_1}\frac{r_3(t,M)}{t+1}\,dt,$$

where $T_1 = 2M(\ln x)^C$,

$$r_3(t,M) = \sum_{M<q_1\leqslant M_1}\sum^{*}_{\chi_1 \bmod q_1}|L(\sigma_1+it,\chi_1)|^{\frac{1}{2}}.$$

From Hölder's inequality and lemma 2 it follows that

$$r_3(t,M) \ll M^2(\ln x)^{\frac{1}{4}(C+1)}.$$

Thus we find

$$R_3 \ll \sqrt{N}(\ln x)^{\frac{3}{2}}(\ln\ln x)^2\sum'_{M\leqslant Q}\frac{1}{M}\,M^2(\ln x)^{\frac{1}{4}(C+1)} \ll$$

$$\ll Q\sqrt{N}(\ln x)^{\frac{1}{4}(C+7)}(\ln\ln x)^2.$$

4°. Estimation of R_4. For given character $\chi \bmod q$ and a zero $\varrho = \beta + i\gamma$ of $L(s,\chi)$ we define the function $g_\chi(\varrho,\sigma)$ as follows:

$$g_\chi(\varrho,\sigma) = \begin{cases} 1, & \text{if } \beta > \sigma,\\ 0, & \text{otherwise.}\end{cases}$$

It's obvious that if $\chi_1 \bmod q_1$ induces $\chi \bmod q$ then $g_{\chi_1}(\varrho,\sigma)$ coincides with $g_\chi(\varrho,\sigma)$ for $\sigma \geqslant \frac{1}{2}$. Thus we have

$$R_4 \leqslant \sum_{q\leqslant Q}\frac{1}{\varphi(q)}\sum_{\substack{\chi \bmod q\\ \chi\neq\chi_0}}\sum_{\substack{|\gamma|\leqslant T(q,\chi)\\ \beta>0.5}}\frac{1}{|\gamma|+1}\int_{0.5}^{1}g_\chi(\varrho,\sigma)N^\sigma|L(\sigma+i\gamma,\chi)|^{\frac{1}{2}}\,d\sigma \leqslant$$

$$\leqslant \int_{0.5}^{1}N^\sigma\sum_{q_1\leqslant Q}\frac{1}{\varphi(q_1)}\sum_{r\leqslant\frac{Q}{q_1}}\frac{\sqrt{\tau(r)}}{\varphi(r)}\sum^{*}_{\chi_1 \bmod q_1}\sum_{\substack{|\gamma|\leqslant T(q,\chi)\\ \beta>0.5}}\frac{g_{\chi_1}(\varrho,\sigma)}{|\gamma|+1}|L(\sigma+i\gamma,\chi_1)|^{\frac{1}{2}}\,d\sigma$$

$$\ll (\ln x)^{\frac{3}{2}}(\ln\ln x)^2 \int_{0.5}^{1} N^{\sigma} \sum_{q_1\leqslant Q}\frac{1}{q_1}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\sum_{\substack{|\gamma|\leqslant T(q,\chi)\\ \beta>0.5}} \frac{g_{\chi_1}(\varrho,\sigma)}{|\gamma|+1}|L(\sigma+i\gamma,\chi_1)|^{\frac12}\,d\sigma$$

$$\ll (\ln x)^{\frac32}(\ln\ln x)^2 \sideset{}{'}\sum_{M\leqslant Q} j(M),$$

where

$$\begin{aligned} j(M) &= \frac{1}{M}\int_{0.5}^{1} N^{\sigma} r_4(\sigma,M)d\sigma,\\ r_4(\sigma,M) &= \sum_{M<q_1\leqslant M_1}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\sum_{\substack{|\gamma|\leqslant T(q,\chi)\\ \beta>0.5}} \frac{g_{\chi_1}(\varrho,\sigma)}{|\gamma|+1}|L(\sigma+i\gamma,\chi_1)|^{\frac12}. \end{aligned}$$

In the below, we consider the cases of «small» and «large» M separately.

4a°. The case of «small» M: $1\leqslant M\leqslant(\ln x)^A$. Obviously we have

$$T(q_1,\chi_1)\leqslant 2M(\ln x)^C\leqslant 2(\ln x)^{A+C}.$$

If $\varrho=\beta+i\gamma$ is a zero of $L(s,\chi_1)$ and $0<|\gamma|\leqslant T(q_1,\chi_1)$ then lemma 4 implies that

$$\beta\leqslant 1-\frac{c_1}{\ln q_1(|\gamma|+2)}\leqslant 1-\frac{c}{\ln\bigl(2(\ln c)^{2A+C}\bigr)}<1-\frac{1}{\sqrt{\ln x}}.$$

Further, if there exists a real zero β of $L(s,\chi_1)$ then lemma 5 implies (with $\varepsilon=(2A)^{-1}$) that

$$\beta\leqslant 1-\frac{c}{q_1^{\varepsilon}}\leqslant 1-\frac{c}{(\ln x)^{A\varepsilon}}=1-\frac{c}{\sqrt{\ln x}}$$

for some $c>0$. Without loss of generality, we may assume that $0<c<1$. Then it follows that for any zero ϱ of the function $L(s,\chi_1)$ under considering the inequality $|\gamma|\leqslant T(q_1,\chi_1)$ implies that $g_{\chi_1}(\varrho,\sigma)=0$ for any $\sigma\geqslant 1-\dfrac{1}{\sqrt{\ln x}}$.

Now, if $\frac12\leqslant\sigma\leqslant 1$ then from lemma 3 it follows that

$$\begin{aligned} |L(\sigma+it,\chi_1)| &\ll \Bigl|\sum_{n\leqslant q_1(|t|+1)}\chi(n)n^{-\sigma-it}\Bigr| + q_1\bigl(q_1(|t|+1)\bigr)^{-\sigma}\\ &\ll \bigl(q_1(|t|+1)\bigr)^{1-\sigma}\ln q_1(|t|+1). \end{aligned}$$

By lemma 7, we easily get

$$|L(\sigma+i\gamma,\chi_1)|^{\frac12}\ll\bigl(q_1T(q_1,\chi_1)\bigr)^{\frac{1-\sigma}{2}}\ln\ln x\ll\bigl(M^2(\ln x)^C\bigr)^{\frac{1-\sigma}{2}}\ln\ln x<$$
$$<(\ln x)^{\frac34 A}\ln\ln x,$$

$$r_4(\sigma,M)\ll(\ln x)^{\frac34 A}\ln\ln x\sum_{M<q\leqslant M_1}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\sum_{|\gamma|\leqslant T(q_1,\chi_1)}\frac{1}{|\gamma|+1}\ll$$
$$<(\ln x)^{\frac34 A}\ln\ln x\sum_{M<q\leqslant M_1}\sideset{}{^*}\sum_{\chi_1 \bmod q_1}\ln^2 q_1T(q_1,\chi_1)\ll(\ln x)^{\frac34 A}(\ln\ln x)^3M^2\ll$$
$$\ll M(\ln x)^{2A}.$$

Thus we finally obtain:

$$j(M) \ll \frac{1}{M}\int_{0.5}^{1-\frac{c}{\sqrt{\ln x}}} N^{\sigma}\, M(\ln x)^{A} d\sigma \ll N^{1-\frac{c}{\sqrt{\ln x}}}(\ln x)^{2A} \ll xe^{-\frac{c}{2}\sqrt{\ln x}}.$$

4b°. The case of «large» M: $(\ln x)^{A} < M \leqslant \sqrt{x}(\ln x)^{-A}$. Let us divide the domain of γ into segments $U < |\gamma| \leqslant U_1$ where

$$U_1 \leqslant 2U, \quad U_1 \leqslant T(q_1,\chi_1) \leqslant 2M(\ln x)^{C} = T_1.$$

Thus we have

$$r_4(\sigma,M) \ll \sum_{U \leqslant T_1}{}' \frac{1}{U} r_4(\sigma,M,U),$$
$$r_4(\sigma,M,U) = \sum_{M<q_1\leqslant M_1} \sum_{\chi_1 \bmod q_1}{}^{*} \sum_{\substack{U<|\gamma|\leqslant U_1\\ \beta>0.5}} g_{\chi_1}(\varrho,\sigma)|L(\sigma+i\gamma,\chi_1)|^{\frac{1}{2}}.$$

Applying Hölder's inequality, we find

$$r_4^4(\sigma,M,U) \ll \Bigl(\sum_{M<q_1\leqslant M_1} \sum_{\chi_1 \bmod q_1}{}^{*} \sum_{\substack{U<|\gamma|\leqslant U_1\\ \beta>0.5}} g_{\chi_1}\Bigr)^3 \times$$
$$\times \Bigl(\sum_{M<q_1\leqslant M_1} \sum_{\chi_1 \bmod q_1}{}^{*} \sum_{\substack{U<|\gamma|\leqslant U_1\\ \beta>0.5}} |L(\sigma+i\gamma,\chi_1)|\Bigr) = r_5 \cdot r_6^3,$$

where the notations r_5, r_6 are obvious.

First we get:

$$r_5 \ll \sum_{M<q_1\leqslant M_1} \sum_{\chi_1 \bmod q_1}{}^{*} \sum_{|\gamma|\leqslant U_1} |L(\sigma+i\gamma,\chi_1)|^2.$$

Let us divide the domain of γ into the segments

$$2n \leqslant \gamma < 2n+1, \quad n = 0, \pm 1, \pm 2, \ldots,$$

and

$$2n+1 \leqslant \gamma < 2(n+1), \quad n = 0, \pm 1, \pm 2, \ldots .$$

Thus the sum r_5 splits into sums r_5' and r_5''. Further, let us sort all the ordinates γ in each segment $2n \leqslant \gamma < 2n+1$ in increasing order:

$$2n \leqslant \gamma^{(1)} \leqslant \gamma^{(2)} \leqslant \ldots \leqslant \gamma^{(s)} < 2n+1.$$

It's obvious that $s = O(\ln q_1(|n|+1))$. Now we place the terms of r_5' that corresponds to the first ordinates $\gamma^{(1)}$ into sum $r_5^{(1)}$. The terms corresponding to the second ordinates $\gamma^{(2)}$ are placed into sum $r_5^{(2)}$ and so on. Thus r_5' splits into $s_0 = O(\ln x)$ sums $r_5^{(s)}$, $s = 1, 2, \ldots, s_0$. The ordinates γ, γ' that corresponds to the neighbouring summands in the sum r_5^{s} satisfy the condition $|\gamma - \gamma'| > 1$.

Finally, we apply the same transformation to the sum r_5''. Thus the sum r_5 splits into $\leqslant 2s_0 = O(\ln x)$ sums r of the following type:

$$r \,=\, \sum_{M<q_1\leqslant M_1}\ \sideset{}{^*}\sum_{\chi_1 \bmod q_1}\ \sideset{}{'}\sum_{|\gamma|\leqslant U} |L(\sigma+i\gamma,\chi_1)|^2$$

where the prime sign means the summation over the «rarefied» ordinates γ.

Taking $Y = M_1(T_1+1)$ we have $Y \geqslant q_1(|\gamma|+1)/\pi$ for all q_1 and γ under considering. Then Lemma 3 implies that

$$r \,\ll\, \sum_{M<q_1\leqslant M_1}\ \sideset{}{^*}\sum_{\chi_1 \bmod q_1}\ \sideset{}{'}\sum_{|\gamma|\leqslant U}\biggl(\Bigl|\sum_{n\leqslant Y}\frac{\chi_1(n)}{n^{\sigma+i\gamma}}\Bigr|^2 + 1\biggr) \,\ll$$

$$\ll\, M^2U \,+\, \sum_{M<q_1\leqslant M_1}\ \sideset{}{^*}\sum_{\chi_1 \bmod q_1}\ \sideset{}{'}\sum_{|\gamma|\leqslant U}\Bigl|\sum_{n\leqslant Y}\frac{\chi_1(n)}{n^{\sigma+i\gamma}}\Bigr|^2.$$

The application of Lemma 1 yields:

$$\sideset{}{'}\sum_{|\gamma|\leqslant U}\Bigl|\sum_{n\leqslant Y}\frac{\chi_1(n)}{n^{\sigma+i\gamma}}\Bigr|^2 \,\leqslant\, j_1 \,+\, 2\sqrt{j_1\, j_2},$$

where

$$j_1 \,=\, \int_{-(U_1+1)}^{U_1+1}\Bigl|\sum_{n\leqslant Y}\frac{\chi_1(n)}{n^{\sigma+it}}\Bigr|^2 dt, \quad j_2 \,=\, \int_{-(U_1+1)}^{U_1+1}\Bigl|\sum_{n\leqslant Y}\frac{\chi_1(n)\ln n}{n^{\sigma+it}}\Bigr|^2 dt.$$

Since $U_1+1 \leqslant 2U+1 \leqslant 3U$, $2\sqrt{j_1j_2} \leqslant j_1+j_2$, by Lemma 2 we obtain

$$r \,\ll\, M^2U \,+\, \sum_{M<q_1\leqslant M_1}\ \sideset{}{^*}\sum_{\chi_1 \bmod q_1}(j_1 \,+\, j_2) \,\ll$$

$$\ll\, M^2U \,+\, \int_{-3U}^{3U}\sum_{M<q_1\leqslant M_1}\ \sideset{}{^*}\sum_{\chi_1 \bmod q_1}\biggl(\Bigl|\sum_{n\leqslant Y}\frac{\chi_1(n)}{n^{\sigma+it}}\Bigr|^2 \,+\, \Bigl|\sum_{n\leqslant Y}\frac{\chi_1(n)\ln n}{n^{\sigma+it}}\Bigr|^2\biggr) dt \,\ll$$

$$\ll\, M^2U \,+\, \int_{-3U}^{3U}\bigl(M^2+Y\bigr)\Bigl(\sum_{n\leqslant Y}\frac{1}{n^{2\sigma}} \,+\, \sum_{n\leqslant Y}\frac{\ln^2 n}{n^{2\sigma}}\Bigr) dt \,\ll$$

$$\ll\, M^2U \,+\, \int_{-3U}^{3U} M^2(\ln x)^{C+3}\, dt \,\ll\, M^2U(\ln x)^{C+3}$$

and therefore

$$r_4(\sigma, M, U) \,\ll\, M^2U(\ln x)^{C+4}.$$

Further, using Lemma 6, we obtain

$$r_6 \leqslant \sum_{q_1\leqslant M_1}\ \sideset{}{^*}\sum_{\chi_1 \bmod q_1}\ \sum_{|\gamma|\leqslant U_1} g_{\chi_1}(\varrho,\sigma) \,=\, \sum_{q_1\leqslant M_1}\ \sideset{}{^*}\sum_{\chi_1 \bmod q_1} N(\sigma; U_1, \chi_1) \,\ll$$

$$\ll\, \bigl(M^2U\bigr)^{\vartheta(\sigma)}(\ln x)^{14}.$$

Therefore,

$$
\begin{aligned}
&r_4^4(\sigma,M,U) \ll M^2U(\ln x)^{C+4}\bigl(M^2U\bigr)^{3\,\vartheta(\sigma)}(\ln x)^{42} \ll \bigl(M^2U\bigr)^{1+3\,\vartheta(\sigma)}(\ln x)^{C+46},\\
&r_4(\sigma,M,U) \ll \bigl(M^2U\bigr)^{\frac{1}{4}+\frac{3}{4}\vartheta(\sigma)}(\ln x)^{\frac{1}{4}C+\frac{23}{2}},\\
&r_4(\sigma,M) \ll \sideset{}{'}\sum_{U\leqslant T_1} U^{-1}\cdot\bigl(M^2U\bigr)^{\frac{1}{4}+\frac{3}{4}\vartheta(\sigma)}(\ln x)^{\frac{1}{4}C+\frac{23}{2}} \ll\\
&\ll (\ln x)^{\frac{1}{4}C+\frac{23}{2}}\sideset{}{'}\sum_{U\leqslant T_1} U^{\frac{3}{4}(\vartheta(\sigma)-1)}M^{\frac{1}{2}+\frac{3}{2}\vartheta(\sigma)}.
\end{aligned}
$$

Since $\vartheta(\sigma)\leqslant 1$ for $\frac{1}{2}\leqslant\sigma\leqslant 1$, it follows that

$$r_4(\sigma,M) \ll M^{\frac{1}{2}+\frac{3}{2}\vartheta(\sigma)}(\ln x)^{\frac{1}{4}C+\frac{25}{2}}$$

and

$$j(M) = M^{-1}\int_{\sigma_1}^{1} N^{\sigma}r_4(\sigma,M)d\sigma \ll (\ln x)^{\frac{1}{4}C+\frac{25}{2}}\int_{0.5}^{1}\psi(\sigma)\,d\sigma,$$

where

$$\psi(\sigma) = x^{\sigma}M^{u(\sigma)},\quad u(\sigma) = -\tfrac{1}{2} + \tfrac{3}{2}\vartheta(\sigma).$$

Let us consider several cases.

i°. For $\frac{1}{2}\leqslant\sigma\leqslant\frac{4}{5}$ we obviously have

$$u(\sigma) = \frac{1}{4} + \frac{3(4-5\sigma)}{4(2-\sigma)} \geqslant \frac{1}{4}.$$

The condition $N\leqslant\sqrt{x}(\ln x)^{-A}$ implies that

$$\psi(\sigma) \leqslant x^{\sigma}\bigl(\sqrt{x}(\ln x)^{-A}\bigr)^{u(\sigma)} \leqslant x^{\sigma+\frac{1}{2}u(\sigma)}(\ln x)^{-\frac{1}{4}A}.$$

Since

$$\sigma + \frac{1}{2}u(\sigma) = 1 - \frac{\bigl(\sigma-\frac{1}{2}\bigr)^2}{2-\sigma} \leqslant 1,$$

we get

$$\psi(\sigma) \leqslant x(\ln x)^{-\frac{1}{4}A}.$$

ii°. Suppose $\frac{4}{5}\leqslant\sigma\leqslant\frac{6}{7}$. Then

$$u(\sigma) = \frac{6-7\sigma}{2\sigma} \geqslant 0$$

and

$$\psi(\sigma) \leqslant x^{\sigma}\bigl(\sqrt{x}(\ln x)^{-A}\bigr)^{u(\sigma)} \leqslant x^{\sigma+\frac{1}{2}u(\sigma)}.$$

From the inequality

$$\frac{d}{d\sigma}\Bigl(\sigma + \tfrac{1}{2}u(\sigma)\Bigr) = 1 - \frac{3}{2\sigma^2} < 0$$

it follows that the function $\sigma + \frac{1}{2}u(\sigma)$ is monotonically decreasing on the segment $\frac{4}{5} \leqslant \sigma \leqslant \frac{6}{7}$ and attains it's maximum at a point $\sigma = \frac{4}{5}$:

$$\sigma \,+\, \tfrac{1}{2}\,u(\sigma) \,\leqslant\, \tfrac{4}{5} \,+\, \tfrac{1}{2}\,u\big(\tfrac{4}{5}\big) \,=\, \tfrac{37}{40}.$$

Thus

$$\psi(\sigma) \,\leqslant\, x^{\frac{37}{40}} \,\leqslant\, x(\ln x)^{-\frac{1}{4}A}.$$

iii°. Suppose $\frac{6}{7} \leqslant \sigma \leqslant \frac{12}{13}$. Then $u(\sigma) \leqslant 0$ and therefore

$$\psi(\sigma) \,\leqslant\, x^{\sigma} \,\leqslant\, x^{\frac{12}{13}} \,\leqslant\, x(\ln x)^{-\frac{1}{4}A}.$$

iv°. Suppose that $\frac{12}{13} \leqslant \sigma \leqslant 1$. Then

$$u(\sigma) \,=\, -\frac{1}{4} \,-\, \frac{13\sigma - 12}{4\sigma} \,\leqslant\, -\frac{1}{4}.$$

Since $M > (\ln x)^{A}$, it implies that

$$\psi(\sigma) \leqslant x^{\sigma}M^{-\frac{1}{4}} \,\leqslant\, x(\ln x)^{-\frac{1}{4}A}.$$

Thus the inequality

$$\psi(\sigma) \,\leqslant\, x(\ln x)^{-\frac{1}{4}A}$$

holds for any σ such that $\frac{1}{2} \leqslant \sigma \leqslant 1$. Finally we get

$$\begin{aligned}
&j(M) \,\ll\, x(\ln x)^{-\frac{1}{4}A + \frac{1}{4}C + \frac{25}{2}},\\
&R_4 \,\ll\, (\ln x)^{\frac{3}{2}}(\ln\ln x)^2\Big(\sideset{}{'}\sum_{M \leqslant (\ln x)^C} xe^{-\frac{c}{2}\sqrt{\ln x}} \,+\, \sideset{}{'}\sum_{(\ln x)^C < M \leqslant Q} x(\ln x)^{-\frac{1}{4}A + \frac{1}{4}C + \frac{25}{2}}\Big)\\
&\;\ll\, x(\ln x)^{-\frac{1}{4}A + \frac{1}{4}C + 15}(\ln\ln x)^2.
\end{aligned}$$

Summing the upper bounds for R_j, $1 \leqslant j \leqslant 4$, we obtain

$$\begin{aligned}
&R \,\ll\, (\ln x)^{\alpha}\Big(N(\ln x)^{-C+3} \,+\, N(\ln x)^{-\frac{3}{4}C + \frac{11}{4}}(\ln\ln x)^2 \,+\\
&+\, Q\sqrt{N}(\ln x)^{\frac{1}{4}(C+1)}(\ln\ln x)^2 \,+\, x(\ln x)^{-\frac{1}{4}A + \frac{1}{4}C + 15}(\ln\ln x)^2\Big) \,\ll\\
&\ll (\ln x)^{\alpha}(\ln\ln x)^2\Big(x(\ln x)^{-C+3} \,+\, x(\ln x)^{-\frac{3}{4}C + \frac{11}{4}} \,+\, x(\ln x)^{-A + \frac{1}{4}(C+1)} \,+\\
&+\, x(\ln x)^{-\frac{1}{4}A + \frac{1}{4}C + 15}(\ln\ln x)^2\Big) \,\ll\\
&\ll\, x(\ln x)^{\alpha}(\ln\ln x)^2\Big((\ln x)^{-\frac{3}{4}C + \frac{11}{4}} \,+\, (\ln x)^{-\frac{1}{4}A + \frac{1}{4}C + 15}\Big).
\end{aligned}$$

Taking $C = \frac{1}{4}(A - 49)$, we get the required inequality:

$$R \,\ll\, x(\ln x)^{\alpha}(\ln\ln x)^2(\ln x)^{-\frac{3}{16}A + \frac{191}{16}} \,\ll\, x(\ln x)^{-\left(\frac{3}{16}A - 12\right)}.$$

If we put $B = \frac{3}{16}A - 12$, then $A = \frac{16}{3}B + 4$. This completes the proof of the lemma.

4. Main theorem

By Lemma 12, it remains to estimate $R_a(x)$ and find an asymptotic formula for $E_a(x)$. Setting $Q = Xd^{-1} = \sqrt{x}d^{-1}(\ln x)^{-A}$, $N = x(d\delta)^{-1}$ in Lemma 13, we obtain

$$R_{a,d,\delta} \ll x(\ln x)^{-\left(\frac{3}{16}A - 12\right)}, \quad R_a(x) \ll x(\ln x)^{-\left(\frac{3}{16}A - 15\right)}.$$

Taking $A = 85\frac{1}{3}$, we get the formula

$$S_a(x) = C(\ln x)\beta(a)E_a(x) + O(x \ln\ln x),$$

where

$$E_a(x) = \sum_{n \leqslant x} \frac{e_a(n)}{\tau(n)}.$$

In order to calculate the sum $E_a(x)$, let us prove that $e_a(n)$ is multiplicative function for any fixed $a \geqslant 1$. Indeed, suppose $(m, n) = 1$. Then there exists a unique decomposition $a = a_1 a_2 a'$ where the factors a_1, a_2, a' are defined as follows. All prime divisors of a_1 and a_2 are among the sets of prime divisors of m and n respectively, and $(a', mn) = 1$. Then

$$(a_1, a_2) = (a_1, a') = (a_2, a') = (a_1, n) = (a_2, m) = 1.$$

Further, if d divides (a, mn) then d has the form $d_1 d_2$ where $d_1|(a, m)$ and $d_2|(a, n)$. Thus,

$$e_a(mn) = \sum_{d|(a,mn)} \frac{\beta\left(\dfrac{an}{d^2}\right)}{\beta(a)} = \sum_{d_1|(a,m)} \sum_{d_2|(a,n)} \frac{\beta\left(\dfrac{a_1 a_2 a' mn}{d_1^2 d_2^2}\right)}{\beta(a_1 a_2 a')} =$$

$$= \sum_{d_1|(a,m)} \sum_{d_2|(a,n)} \frac{\beta\left(\dfrac{a_1 m}{d_1^2} \cdot \dfrac{a_2 n}{d_2^2}\right)\beta(a')}{\beta(a_1 a_2)\beta(a')}.$$

Since the numbers $\dfrac{a_1 m}{d_1^2}$ and $\dfrac{a_2 n}{d_2^2}$ are integral and coprime, we have

$$e_a(mn) = \sum_{d_1|(a_1,m)} \frac{\beta\left(\dfrac{a_1 m}{d_1^2}\right)}{\beta(a_1)} \sum_{d_2|(a_2,n)} \frac{\beta\left(\dfrac{a_2 n}{d_2^2}\right)}{\beta(a_2)} =$$

$$= \sum_{d_1|(a_1,m)} \frac{\beta\left(\dfrac{a_1 m}{d_1^2}\right)\beta(a_2 a')}{\beta(a_1)\beta(a_2 a')} \sum_{d_2|(a_2,n)} \frac{\beta\left(\dfrac{a_2 n}{d_2^2}\right)\beta(a_1 a')}{\beta(a_2)\beta(a_1 a')} =$$

$$= \sum_{d_1|(a_1,m)} \frac{\beta\left(\dfrac{am}{d_1^2}\right)}{\beta(a)} \sum_{d_2|(a_2,n)} \frac{\beta\left(\dfrac{an}{d_2^2}\right)}{\beta(a)} = \sum_{d_1|(a,m)} \frac{\beta\left(\dfrac{am}{d_1^2}\right)}{\beta(a)} \sum_{d_2|(a,n)} \frac{\beta\left(\dfrac{an}{d_2^2}\right)}{\beta(a)} =$$

$$= e_a(m)e_a(n).$$

Suppose $\operatorname{Re} s > 1$. By definition, put

$$F_a(s) \,=\, \sum_{n=1}^{+\infty} \frac{e_a(n)}{\tau(n)}\, n^{-s}.$$

Then

$$F_a(s) \,=\, \prod_p F_{a,p}(s), \quad F_{a,p}(s) \,=\, 1 \,+\, \sum_{k=1}^{+\infty} \frac{e_a(p^k)}{k+1}\, p^{-ks}.$$

Since

$$e_a(n) \,=\, \frac{\beta(an)}{\beta(a)} \,=\, \beta(n)$$

for the case $(a, n) = 1$, we obtain

$$\begin{aligned} F_a(s) \,&=\, \prod_{p \nmid a}\Big(1 \,+\, \sum_{k=1}^{+\infty} \frac{\beta(p^k)}{k+1}\, p^{-ks}\Big) \prod_{p \mid a} F_{a,p}(s) \,= \\ &= \prod_{p \nmid a}\Big(1 \,+\, \beta(p) \sum_{k=1}^{+\infty} \frac{p^{-ks}}{k+1}\Big) \prod_{p \mid a} F_{a,p}(s) \,= \\ &= \prod_{p \nmid a}\Big(1 - \beta(p) - \beta(p) p^s \ln(1 - p^{-s})\Big) \prod_{p \mid a} F_{a,p}(s) \,=\, F(s)\psi_a(s), \end{aligned}$$

where

$$F(s) \,=\, \prod_p \Big(1 - \beta(p) - \beta(p) p^s \ln(1 - p^{-s})\Big),$$

$$\psi_a(s) \,=\, \prod_{p \mid a} \frac{1 \,+\, \displaystyle\sum_{k=1}^{+\infty} \frac{e_a(p^k)}{k+1}\, p^{-ks}}{1 - \beta(p) - \beta(p) p^s \ln(1 - p^{-s})}.$$

Further, $F(s) = \sqrt{\zeta(s)}\, \Phi(s)$, where

$$\Phi(s) \,=\, \prod_p \Phi_p(s), \quad \Phi_p(s) \,=\, \Big(1 - \frac{1}{p^s}\Big)^{\frac{1}{2}} \big(1 - \beta(p) - \beta(p) p^s \ln(1 - p^{-s})\big).$$

Setting

$$1 - \beta(p) - \beta(p) p^s \ln(1 - p^{-s}) \,=\, 1 + \frac{1}{2p^s} + u(s), \quad \Big(1 - \frac{1}{p^s}\Big)^{\frac{1}{2}} \,=\, 1 - \frac{1}{2p^s} + v(s)$$

and using the decomposition

$$\begin{aligned} \beta(p) \,&=\, 1 \,-\, \frac{p}{p^2 - p + 1} \,=\, 1 - \frac{p(p+1)}{p^3 + 1} \,=\, 1 \,-\, \frac{1}{p}\Big(1 \,+\, \frac{1}{p}\Big)\Big(1 \,+\, \frac{1}{p^3}\Big)^{-1} \,= \\ &= \,1 \,+\, \sum_{k=0}^{+\infty} (-1)^{k+1} \Big(\frac{1}{p^{3k+1}} \,+\, \frac{1}{p^{3k+2}}\Big), \end{aligned}$$

we find

$$\begin{aligned} u(s) &= \frac{1}{3p^{2s}} + \frac{1}{4p^{3s}} + \frac{1}{5p^{4s}} + \ldots - \\ &- \left(\frac{1}{p} + \frac{1}{p^2} - \frac{1}{p^4} - \frac{1}{p^5} + \ldots\right)\left(\frac{1}{2p^s} + \frac{1}{3p^{2s}} + \frac{1}{4p^{3s}} + \ldots\right), \\ v(s) &= \sum_{k=2}^{+\infty} 2^{-2k}\binom{2k}{k}\frac{p^{-ks}}{2k-1}. \end{aligned}$$

Suppose now $\sigma > \frac{1}{2}$. Then

$$\begin{aligned} &|u(s)| \leqslant \\ &\leqslant \frac{1}{3p^{2\sigma}} + \frac{1}{4p^{3\sigma}} + \frac{1}{5p^{4\sigma}} + \ldots + \left(\frac{1}{p} + \frac{1}{p^2}\right)\left(\frac{1}{2p^{\sigma}} + \frac{1}{3p^{2\sigma}} + \frac{1}{4p^{3\sigma}} + \ldots\right) \leqslant \\ &\leqslant \frac{1}{3p^{2\sigma}}\left(1 + \frac{1}{p^{\sigma}} + \frac{1}{p^{2\sigma}} + \ldots\right) + \frac{3}{2p}\cdot\frac{1}{2p^{\sigma}}\left(1 + \frac{1}{p^{\sigma}} + \frac{1}{p^{2\sigma}} + \ldots\right) \leqslant \\ &\leqslant \left(\frac{1}{3p^{2\sigma}} + \frac{3}{4p^{\sigma+1}}\right)\frac{1}{1-p^{-\sigma}} \leqslant \frac{\sqrt{2}}{\sqrt{2}-1}\left(\frac{1}{3p^{2\sigma}} + \frac{3}{4p^{\sigma+1}}\right), \end{aligned}$$

and therefore

$$|u(s)| \leqslant \frac{\sqrt{2}}{\sqrt{2}-1}\left(\frac{1}{3} + \frac{3}{4}\right)\max\left(\frac{1}{p^{2\sigma}}, \frac{1}{p^{\sigma+1}}\right) \leqslant \begin{cases} 3.7p^{-2\sigma}, & \text{if } \frac{1}{2} < \sigma \leqslant 1, \\ 3.7p^{-(\sigma+1)}, & \text{if } \sigma \geqslant 1. \end{cases}$$

Further,

$$\begin{aligned} |v(s)| &\leqslant \sum_{k=2}^{+\infty} 2^{-2k}\binom{2k}{k}\frac{p^{-k\sigma}}{2k-1} = 1 - \frac{1}{2p^{\sigma}} - \sqrt{1 - \frac{1}{p^{\sigma}}} = \\ &= \frac{\left(1 - \frac{1}{2p^{\sigma}} - \sqrt{1 - p^{-\sigma}}\right)\left(1 - \frac{1}{2p^{\sigma}} + \sqrt{1 - p^{-\sigma}}\right)}{\left(1 - \frac{1}{2p^{\sigma}} + \sqrt{1 - p^{-\sigma}}\right)} = \\ &= \frac{\left(1 - \frac{1}{2p^{\sigma}}\right)^2 - \left(1 - \frac{1}{p^{\sigma}}\right)}{\left(1 - \frac{1}{2p^{\sigma}}\right) + \sqrt{1 - p^{-\sigma}}} \leqslant \frac{\frac{1}{4p^{2\sigma}}}{1 - \frac{1}{2\sqrt{2}} + \sqrt{1 - \frac{1}{\sqrt{2}}}} < \frac{1}{4p^{2\sigma}} \end{aligned}$$

for any $\sigma > \frac{1}{2}$. In particular, in the case $\sigma \geqslant 1$ we get

$$|v(s)| < \frac{1}{4p^{2\sigma}} \leqslant \frac{1}{4p^{\sigma+1}}.$$

Thus we obtain for $\frac{1}{2} \leqslant \sigma \leqslant 1$

$$|\Phi_p(s)| = \left|1 - \frac{1}{4p^{2s}} + u(s) - v(s) - \frac{u(s)}{2p^s} - \frac{v(s)}{2p^s} - u(s)v(s)\right| \leqslant$$

$$\leqslant 1 + \frac{1}{4p^{2\sigma}} + \frac{3.7}{p^{2\sigma}} + \frac{1}{4p^{2\sigma}} + \frac{1}{2p^{\sigma}}\Big(\frac{3.7}{p^{2\sigma}} + \frac{1}{4p^{2\sigma}}\Big) + \frac{3.7}{4p^{4\sigma}} <$$
$$< 1 + \frac{1}{2p^{2\sigma}} + \frac{3.7}{p^{2\sigma}} + \frac{3.95}{2\sqrt{2}p^{2\sigma}} + \frac{3.7}{8p^{2\sigma}} < 1 + \frac{7}{p^{2\sigma}} < \Big(1 - \frac{1}{p^{2\sigma}}\Big)^{7}$$

and therefore

$$|\Phi(s)| < \prod_p \Big(1 - \frac{1}{p^{2\sigma}}\Big)^{7} = \zeta^7(2\sigma) < \Big(\frac{2\sigma}{2\sigma - 1}\Big)^{7} = \big(\sigma - \tfrac{1}{2}\big)^{-7}.$$

Similarly, in the case $\sigma \geqslant 1$ we have

$$|\Phi(s)| < \zeta^7(\sigma + 1) \leqslant \zeta^7(2) = O(1).$$

Thus we get for $\operatorname{Re} s > 1$:

$$F_a(s) = \sqrt{\zeta(s)}\,\Phi_a(s), \quad \Phi_a(s) = F(s)\psi_a(s)$$

where the function $\Phi_a(s)$ is regular in the half-plane $\operatorname{Re} s > \frac{1}{2}$ and obeys the inequality

$$|\Phi_a(s)| \ll \max\Big(\big(\sigma - \tfrac{1}{2}\big)^{-7}, 1\Big).$$

Now Lemma 8 implies that

$$E_a(x) = \frac{x}{\sqrt{\ln x}}\Big(\frac{\Phi_a(1)}{\sqrt{\pi}} + O\Big(\frac{1}{\ln x}\Big)\Big)$$

and therefore

$$S_a(x) = K(a)x\sqrt{\ln x} + O(x \ln\ln x)$$

where

$$K(a) = \frac{1}{\sqrt{\pi}}C\beta(a)\Phi_a(1).$$

Finally, considering the constant $K(a)$, we get $K(a) = K \cdot \kappa(a)$, where

$$K = \frac{1}{\sqrt{\pi}}C\Phi(1) = \frac{C}{\sqrt{\pi}}\prod_p \sqrt{1 - \frac{1}{p}}\,\big(1 - \beta(p) - \beta(p)p\ln(1 - p^{-1})\big) =$$
$$= \frac{1}{\sqrt{\pi}}\prod_p \Big(1 + \frac{1}{p(p-1)}\Big)\sqrt{1 - \frac{1}{p}}\,\frac{p\big(1 + (p-1)^2\ln\frac{p}{p-1}\big)}{p^2 - p + 1} =$$
$$= \frac{1}{\sqrt{\pi}}\prod_p \sqrt{1 - \frac{1}{p}}\,\frac{p^2 - p + 1}{p(p-1)}\,\frac{p\big(1 + (p-1)^2\ln\frac{p}{p-1}\big)}{p^2 - p + 1} =$$
$$= \frac{1}{\sqrt{\pi}}\prod_p \Big(\frac{1}{\sqrt{p(p-1)}} + \sqrt{1 - \frac{1}{p}}\,(p-1)\ln\frac{p}{p-1}\Big),$$

$$\kappa(a) = \beta(a)\psi_a(1) = \beta(a)\prod_{p|a}\frac{1 + \sum\limits_{k=1}^{+\infty}\frac{e_a(p^k)}{k+1}\,p^{-k}}{1 + \beta(p)\sum\limits_{k=1}^{+\infty}\frac{p^{-k}}{k+1}}.$$

Theorem is completely proved.

Remark. The above arguments may be applied to the sums of the following type:

$$C_a = \sum_{a<n\leqslant x} \frac{\tau(n)}{\tau(n-a)}, \quad S_{a,k}(x) = \sum_{n\leqslant x} \frac{\tau(n)}{\tau_k(n+a)},$$

where $a \geqslant 1$ and $\tau_k(n)$ denotes the number of solutions of the equation

$$x_1 \ldots x_k = n$$

in the natural numbers $x_1, \ldots, x_k$. One can obtain the formula

$$S_{a,k}(x) = K(a,k)\, x(\ln x)^{\frac{1}{k}} + O(x \ln\ln x).$$

5. Calculation of the constants $\kappa(a)$

First we note that $\kappa(a)$ is the multiplicative function of a. Indeed, suppose $a = a_1a_2$ where $(a_1, a_2) = 1$, and let p be a prime divisor of a_1. Then $(p, a_2) = 1$ and for any $k \geqslant 1$

$$e_a(p^k) = \frac{1}{\beta(a)} \sum_{d|(a,p^k)} \beta\Big(\frac{ap^k}{d^2}\Big) = \frac{1}{\beta(a_1)\beta_{a_2}} \sum_{d|(a_1,p^k)} \beta\Big(a_2 \cdot \frac{a_1p^k}{d^2}\Big) =$$
$$= \frac{1}{\beta(a_1)} \sum_{d|(a_1,p^k)} \beta\Big(\frac{a_1p^k}{d^2}\Big) = e_{a_1}(p^k).$$

Now it follows that

$$\kappa(a_1a_2) =$$
$$= \beta(a_1a_2) \prod_{p|a_1} \frac{1 + \sum\limits_{k=1}^{+\infty} \dfrac{e_a(p^k)}{k+1}\, p^{-k}}{1 - \beta(p) - \beta(p)p\ln\dfrac{p}{p-1}} \cdot \prod_{p|a_2} \frac{1 + \sum\limits_{k=1}^{+\infty} \dfrac{e_a(p^k)}{k+1}\, p^{-k}}{1 - \beta(p) - \beta(p)p\ln\dfrac{p}{p-1}} =$$
$$= \beta_{a_1} \prod_{p|a_1} \frac{1 + \sum\limits_{k=1}^{+\infty} \dfrac{e_{a_1}(p^k)}{k+1}\, p^{-k}}{1 - \beta(p) - \beta(p)p\ln\dfrac{p}{p-1}} \cdot \beta_{a_2} \prod_{p|a_2} \frac{1 + \sum\limits_{k=1}^{+\infty} \dfrac{e_{a_2}(p^k)}{k+1}\, p^{-k}}{1 - \beta(p) - \beta(p)p\ln\dfrac{p}{p-1}} =$$
$$= \kappa(a_1)\kappa(a_2).$$

Further, $e_a(p^k) = \beta(p)$ for $(a,p) = 1$ and

$$e_a(p^k) = \begin{cases} \min(k+1, m+1), & \text{if } k \neq m, \\ m + \dfrac{1}{\beta(p)}, & \text{if } k = m, \end{cases}$$

for the case $a = p^m a_1$, $(a_1, p) = 1$, $m \geqslant 1$. Now we calculate the values of $\kappa(a)$ for the cases $a = p, p^2, p^3, p^4$ (p is prime).

1°. Suppose $a = p$; then

$$e_a(p) \;=\; 1 \;+\; \frac{1}{\beta(p)}, \quad e_a(p^k) \;=\; 2, \quad k = 2, 3, 4, \dots\;.$$

Therefore,

$$1 \;+\; \sum_{k=1}^{+\infty} \frac{e_a(p^k)}{k+1}\, p^{-k} \;=\; 1 \;+\; \frac{1}{2p}\Big(1 \;+\; \frac{1}{\beta(p)}\Big) \;+\; 2\sum_{k=2}^{+\infty} \frac{p^{-k}}{k+1} \;=$$

$$=\; 2p \ln \frac{p}{p-1} \;-\; 1 \;-\; \frac{1}{2p} \;+\; \frac{1}{2p\beta(p)}.$$

Hence,

$$\kappa(p) \;=\; \frac{2p \ln \dfrac{p}{p-1} \;-\; 1 \;-\; \dfrac{1}{2p} \;+\; \dfrac{1}{2p\beta(p)}}{p \ln \dfrac{p}{p-1} - 1 + \dfrac{1}{\beta(p)}}.$$

In particular,

$$\kappa(2) \;=\; \frac{2 \ln 2 \;-\; \frac{1}{4}}{\ln 2 + 1} \;=\; 0.671\,113\,754\,\dots$$

$$\kappa(3) \;=\; \frac{2 \ln \frac{3}{2} \;-\; \frac{7}{24}}{\ln \frac{3}{2} + \frac{1}{4}} \;=\; 0.792\,206\,241\,\dots$$

$$\kappa(5) \;=\; \frac{2 \ln \frac{5}{4} \;-\; \frac{31}{160}}{\ln \frac{5}{4} + \frac{1}{16}} \;=\; 0.884\,098\,735\,\dots$$

$$\kappa(7) \;=\; \frac{2 \ln \frac{7}{6} \;-\; \frac{71}{504}}{\ln \frac{7}{6} + \frac{1}{36}} \;=\; 0.920\,297\,714\,\dots$$

$$\kappa(11) \;=\; \frac{2 \ln \frac{11}{10} \;-\; \frac{199}{2200}}{\ln \frac{11}{10} + \frac{1}{100}} \;=\; 0.951\,150\,347\,\dots$$

$$\kappa(13) \;=\; \frac{2 \ln \frac{13}{12} \;-\; \frac{287}{3744}}{\ln \frac{13}{12} + \frac{1}{144}} \;=\; 0.959\,100\,63\,\dots$$

$$\kappa(17) \;=\; \frac{2 \ln \frac{17}{16} \;-\; \frac{511}{8704}}{\ln \frac{17}{16} + \frac{1}{256}} \;=\; 0.969\,157\,895\,\dots$$

$$\kappa(19) \;=\; \frac{2 \ln \frac{19}{18} \;-\; \frac{647}{12312}}{\ln \frac{19}{18} + \frac{1}{324}} \;=\; 0.972\,537\,955\,\dots\;.$$

2°. Suppose $a = p^2$; then

$$1 \;+\; \sum_{k=1}^{+\infty} \frac{e_a(p^k)}{k+1}\, p^{-k} \;=\; 3p \ln \frac{p}{p-1} \;-\; 2 \;-\; \frac{1}{2p} \;-\; \frac{1}{3p^2} \;+\; \frac{1}{3p^2\beta(p)},$$

$$\kappa(p^2) \;=\; \frac{3p \ln \dfrac{p}{p-1} \;-\; 2 \;-\; \dfrac{1}{2p} \;-\; \dfrac{1}{3p^2} \;+\; \dfrac{1}{3p^2\beta(p)}}{p \ln \dfrac{p}{p-1} - 1 + \dfrac{1}{\beta(p)}}.$$

In particular,

$$\begin{aligned}\kappa(2^2) &= \frac{3\ln 2 \;-\; \frac{25}{24}}{\ln 2 + 1} = 0.612\,926\,558\,\ldots\\ \kappa(3^2) &= \frac{3\ln \frac{3}{2} \;-\; \frac{77}{108}}{\ln \frac{3}{2} + \frac{1}{4}} = 0.768\,053\,638\,\ldots\,.\end{aligned}$$

3°. Suppose $a = p^3$; then

$$1 + \sum_{k=1}^{+\infty} \frac{e_a(p^k)}{k+1}\, p^{-k} = 4p\ln\frac{p}{p-1} - 3 - \frac{1}{p} - \frac{1}{3p^2} - \frac{1}{4p^3} + \frac{1}{4p^3\beta(p)},$$

$$\kappa(p^3) = \frac{4p\ln\dfrac{p}{p-1} - 3 - \dfrac{1}{p} - \dfrac{1}{3p^2} - \dfrac{1}{4p^3} + \dfrac{1}{4p^3\beta(p)}}{p\ln\dfrac{p}{p-1} - 1 + \dfrac{1}{\beta(p)}}.$$

In particular,

$$\kappa(2^3) = \frac{4\ln 2 \;-\; \frac{169}{96}}{\ln 2 + 1} = 0.597\,805\,121\,\ldots\,.$$

4°. Suppose $a = p^4$; then

$$1 + \sum_{k=1}^{+\infty} \frac{e_a(p^k)}{k+1}\, p^{-k} = 5p\ln\frac{p}{p-1} - 4 - \frac{3}{2p} - \frac{2}{3p^2} - \frac{1}{4p^3} - \frac{1}{5p^4} + \frac{1}{5p^4\beta(p)},$$

$$\kappa(p^3) = \frac{5p\ln\dfrac{p}{p-1} - 4 - \dfrac{3}{2p} - \dfrac{2}{3p^2} - \dfrac{1}{4p^3} - \dfrac{1}{5p^4} + \dfrac{1}{5p^4\beta(p)}}{p\ln\dfrac{p}{p-1} - 1 + \dfrac{1}{\beta(p)}}.$$

In particular,

$$\kappa(2^4) = \frac{5\ln 2 \;-\; \frac{2363}{960}}{\ln 2 + 1} = 0.593\,142\,51\,\ldots\,.$$

BIBLIOGRAPHY

[1] A.A. Karatsuba, *Basic analytic number theory*, Springer-Verlag, Berlin 1993.

[2] H. Davenport, *Multiplicative number theory*, Markham publishing company, Chicago 1967.

[3] H. Montgomery, *Topics in multiplicative number theory. Lecture notes in mathematics. 227*, Springer-Verlag, Berlin - Heidelberg - New York 1971.